



Worthington purifies trypsin from bovine pancreas. Our enzyme is extensively used in protein sequencing work and for tissue dissociation. For protein sequencing, the Worthington product Code: TRTPCK has been used successfully for many years. Worthington now offers two 'Sequencing Grades' which are purified and characterized to meet the most stringent sequencing application requirements.

Likewise, for tissue culture work, Worthington trypsin has been used by many researchers through the years. We do not offer crude grades such as NF 1:250. Difficulties are often encountered when using these crude preparations including incomplete solubility, lot-to-lot variability, and cell toxicity. While purified trypsin alone may have limited effectiveness for tissue dissociation since it shows little reactivity with extracellular proteins, combinations of purified trypsin and other enzymes such as collagenase and elastase have proven effective for dissociation. The purified enzyme is especially useful for cell harvesting by a process called "trypsinization".

For applications requiring particularly clean trypsin, Worthington offers VMF (Virus and Mycoplasma Free) Trypsin which is a specially processed trypsin. The product is irradiated to eliminate microorganisms and is quality control tested specifically against Bovine Diarrhea virus, Infectious Bovine Rhinotracheitis (IBR) virus, Herpes virus, Enterovirus, Adenoviruses, Bovine syncytial virus, Parainfluenza type 3 (PI3-SF) and Parvovirus. Additionally, the material is tested to be free of mycoplasma. As a result of the purification process, extraneous proteases and nucleases which could be harmful to cell lines are removed. The trypsin specific activity is very high allowing lower working concentration (0.01% range) resulting in less cell damage. An irradiated form of our TPCK-treated trypsin is also available. See product code TRTVMF.

| Description | Activity | Code | Catalog No. | Size | Price |
|---|---|----------------|-------------|----------|----------|
| SequENZ® Trypsin, Modified, Sequencing Grade Worthington TPCK-treated trypsin, code TRTPCK, chemically modified to reduce autolysis and increase stability while retaining its specificity. Supplied as a lyophilized powder. Exhibits a single band on SDS-PAGE. Store at -20°C. PROTECT FROM MOISTURE. REQUIRES SPECIAL SHIPPING: ICE PACK. | ≥ 150 Units per mg protein (≥ 8,625 BAAE/2875 USP/NF units per mg protein) | TRSEQZ | LS02120 | 4x25 µg | \$ 82.00 |
| | | | LS02122 | 4x100 µg | 227.00 |
| | | | LS02123 | 1 mg | 422.00 |
| | | | LS02124 | Bulk | Inquire |
| Trypsin, Purified Sequencing Grade II Trypsin, treated with L-(tosylamido 2-phenyl) ethyl chloromethyl ketone (TPCK) to inhibit contaminant chymotryptic activity according to Kostka and Carpenter, <i>J. Biol. Chem.</i> , 239, 1799, (1964). Supplied as a lyophilized powder. Store at -20°C. PROTECT FROM MOISTURE. REQUIRES SPECIAL SHIPPING: ICE PACK. | ≥ 150 Units per mg protein (≥ 8,625 BAAE/2875 USP/NF units per mg protein) | TRSEQII | LS02115 | 4x25 µg | \$ 70.00 |
| | | | LS02117 | 4x100 µg | 191.00 |
| | | | LS02119 | 1 mg | 355.00 |
| | | | LS02118 | Bulk | Inquire |
| Trypsin, TPCK Treated Treated with L-(tosylamido 2-phenyl) ethyl chloromethyl ketone (TPCK) to inhibit contaminant chymotryptic activity according to Kostka and Carpenter, <i>J. Biol. Chem.</i> , 239, 1799, (1964). Dialyzed against 1mM HCl and lyophilized. Store at 2 - 8°C. PROTECT FROM MOISTURE. | ≥ 180 Units per mg protein (≥ 10,350 BAAE/3,450 USP/NF u/mg protein) | TRTPCK | LS003740 | 100 mg | \$ 95.00 |
| | | | LS003741 | 500 mg | 335.00 |
| | | | LS003744 | 1 gm | 570.00 |
| | | | LS003742 | Bulk | Inquire |

| Description | Activity | Code | Catalog No. | Size | Price |
|--|--|---------------|-------------|----------|-----------|
| Trypsin 3X Supplied as a chromatographically purified, diafiltered and lyophilized powder. Store at 2 - 8°C. PROTECT FROM MOISTURE. | ≥ 180 Units per mg protein (≥ 10,350 BAAE/3,450 USP/NF u/mg protein) | TRL3 | LS003708 | 100 mg | \$ 45.00 |
| | | | LS003707 | 1 gm | 266.00 |
| | | | LS003709 | Bulk | Inquire |
| Trypsin 2X Supplied as a dialyzed and lyophilized powder. Store at 2 - 8°C. PROTECT FROM MOISTURE. | ≥ 180 Units per mg protein (≥ 10,350 BAAE/3,450 USP/NF u/mg protein) | TRL | LS003702 | 100 mg | \$ 24.00 |
| | | | LS003703 | 1 gm | 134.00 |
| | | | LS003704 | 10 gm | 970.00 |
| | | | LS003706 | Bulk | Inquire |
| Trypsin, 0.22µ Filtered Trypsin chromatographically purified, diafiltered, (Code TRL3) filtered through a 0.22 micron pore size membrane and lyophilized in sterile vials. This product is not tested for pyrogenicity. Store at 2 - 8°C. PROTECT FROM MOISTURE. | ≥ 180 Units per mg protein (≥ 10,350 BAAE/3,450 USP/NF u/mg protein) | TRLS | LS003736 | 50 mg | \$ 42.00 |
| | | | LS003734 | 5x50 mg | 180.00 |
| | | | LS003738 | Bulk | Inquire |
| Trypsin, Sterile, Irradiated Free of virus and mycoplasma. Chromatographically purified. Store at 2 - 8°C. | ≥ 180 Units per mg protein (≥ 10,350 BAAE/3,450 USP/NF u/mg protein) | TRLVMF | LS004454 | 100 mg | \$ 95.00 |
| | | | LS004452 | 5x100 mg | 349.00 |
| Trypsin, TPCK-Treated, Irradiated Chromatographically purified trypsin treated with L-(tosylamido-2-phenyl) ethyl chloromethyl ketone (TPCK) to inhibit contaminating chymotryptic activity according to (Kostka and Carpenter, <i>J. Biol. Chem.</i> 239, 1799, 1964), Code: TRTPCK, lyophilized, irradiated and tested for the absence of mycoplasma and extraneous virus according to 9 CFR 113.53c. Each vial is filled to contain ≥ 100 mg. Store at 2-8°C. PROTECT FROM MOISTURE. | ≥ 180 Units per mg protein (≥ 10,350 BAAE/3,450 USP/NF u/mg protein) | TRTVMF | LS003750 | 100 mg | \$ 140.00 |
| | | | LS003752 | 5x100 mg | 536.00 |

Trypsin is a pancreatic serine protease with substrate specificity based upon positively charged lysine and arginine side chains. The molecular weight of trypsinogen is 24,000 daltons and 23,800 daltons for trypsin. The optimum pH is 8.0. Trypsin is inhibited by organophosphorus compounds such as diisopropyl fluorophosphate (DFP) and natural inhibitors from pancreas. Soybean, lima bean, and egg white are also sources of natural inhibitors.

Stability: Most grades of Worthington trypsin are stable for 2 - 3 years when stored at 2 - 8°C.

Storage: Store at 2 - 8°C. Protect from moisture.

Unit Definition: 1 unit hydrolyzes 1 µmole of p-toluene-sulfonyl-L-arginine methyl ester (TAME) per minute at 25°C, pH 8.2, in the presence of 10mM calcium ion. 1mg trypsin ≥180 TAME units, 10,350 BAAE units, 3,450 USP/NF units.

Technical Notes

The virus and mycoplasma free trypsins (Codes: TRLVMF/TRTVMF) have been subjected to gamma irradiation and filtered through 22µm pore size membrane and tested for bioburden. One TAME unit equals 19.17 NF/USP (BAAE) units or 57.5 BAAE units.

Related Products

Cell Isolation Optimizing System • Chymotrypsin • Clostripain (Endoproteinase-Arg-C) • Collagenase • Deoxyribonuclease I
Hepatocyte Isolation System • Hyaluronidase • Neonatal Cardiomyocyte Isolation System • Neutral Protease (Dispase®) • Papain Dissociation System
Protease Staph (Endoproteinase-Glu-C) • Proteinase K • *STEMxyme*® 1 & 2 Collagenase/Neutral Protease Blends • Trypsin Inhibitors