

## Assessing the Metabolic Competence of Sandwich-Cultured Mouse Primary Hepatocytes<sup>S</sup>

Karen Mathijs, Anne S. Kienhuis, Karen J. J. Brauers, Danyel G. J. Jennen, Agustin Lahoz, Jos C. S. Kleinjans, and Joost H. M. van Delft

Department of Health Risk Analysis and Toxicology, Maastricht University, Maastricht, The Netherlands (K.M., A.S.K., K.J.J.B., D.G.J.J., J.C.S.K., J.H.M.v.D.); Netherlands Toxicogenomics Center, Maastricht University, Maastricht, The Netherlands (K.M., D.G.J.J., J.C.S.K., J.H.M.v.D.); and Unidad Mixta Hospital La Fe-Advancell, Centro de Investigación Hospital La Fe, Valencia, Spain (A.L.)

Received November 20, 2008; accepted February 26, 2009

### ABSTRACT:

Primary human and rat hepatocyte cultures are well established in vitro systems used in toxicological studies. However, whereas transgenic mouse models provide an opportunity for studying mechanisms of toxicity, mouse primary hepatocyte cultures are less well described. The potential usefulness of a mouse hepatocyte-based in vitro model was assessed in this study by investigating time-dependent competence for xenobiotic metabolism and gene expression profiles. Primary mouse hepatocytes, isolated using two-step collagenase perfusion, were cultured in a collagen sandwich configuration. Gene expression profiles and the activities of various cytochrome P450 (P450) enzymes were determined after 0, 42, and 90 h in culture. Principal component analysis of gene expression profiles shows that replicates per time point

are similar. Gene expression levels of most phase I biotransformation enzymes decrease to approximately 69 and 57% of the original levels at 42 and 90 h, respectively, whereas enzyme activities for most of the studied P450s decrease to 59 and 34%. The decrease for phase II gene expression is only to 96 and 92% of the original levels at 42 and 90 h, respectively. Pathway analysis reveals initial effects at the level of proteins, external signaling pathways, and energy production. Later effects are observed for transcription, translation, membranes, and cell cycle-related gene sets. These results indicate that the sandwich-cultured primary mouse hepatocyte system is robust and seems to maintain its metabolic competence better than that of the rat hepatocyte system.

Toxicological studies designed to assess safety and possible toxicity of compounds in human populations predominantly rely on the use of animal systems. Because animal experiments are expensive, time-consuming, and connected with ethical drawbacks, numerous attempts have been made to find a reliable way of predicting in vivo toxicity in humans by means of in vitro models in recent years.

Traditionally, because the liver plays a major role in the metabolism of many compounds and also represents an important target organ in systemic toxicity, hepatic models are frequently used as an in vitro alternative in pharmacological, toxicological, and metabolic studies (Blaauboer et al., 1994; Davila et al., 1998). Hepatic cell lines, primary hepatocyte cultures, and precision-cut liver slices from various species are well established in vitro systems for these studies (Blaauboer et al., 1994; Davila et al., 1998; Schaeffner et al., 2005). As precision-cut liver slices have a very short live span and immor-

talized cell lines have lost many liver-specific functions, primary cultures of hepatocytes are preferentially used for in vitro studies on liver toxicity (Berry et al., 1997; Richert et al., 2002; Schaeffner et al., 2005).

Primary mammalian hepatocytes largely retain their liver-specific functions when freshly derived from the animal. However, long-term cultures of functional hepatocytes are difficult to establish. To increase the longevity and maintain differentiated functions of hepatocytes in primary cultures, cells are cultured in a sandwich configuration of collagen-collagen and in serum-free culture medium (Tuschl and Mueller, 2006). Hepatocytes cultured in a sandwich configuration reorganize to form an architecture similar to that found in the liver and are able to form functional bile canaliculi networks and gap junctions. In such a sandwich configuration, hepatocytes can be cultured for a longer period compared with cultures on single layers of collagen (LeCluyse et al., 2000a,b; Boess et al., 2003; Kienhuis et al., 2007).

The use of primary human hepatocytes is preferential to predict in vivo toxicity in humans but is hampered by the limited availability of donor material and the large variability between the donors (Schaeffner et al., 2005). Therefore, primary hepatocytes isolated from other mammals are used as an alternative to human tissue. However, the rapid decline in liver-specific functions, in particular cytochrome

This work was supported in part by The Netherlands Genomics Initiative; The Netherlands Organisation for Scientific Research; and the carcinoGENOMICS FP6 project sponsored by the European Union [Grant PL037712].

Article, publication date, and citation information can be found at <http://dmd.aspetjournals.org>.

doi:10.1124/dmd.108.025775.

<sup>S</sup> The online version of this article (available at <http://dmd.aspetjournals.org>) contains supplemental material.

**ABBREVIATIONS:** P450, cytochrome P450; DMSO, dimethylsulfoxide; SAM, significance analysis of microarrays; PCA, principal component analysis.

P450 (P450) enzyme activity, in rat liver in vitro systems limits the use of these in studies for testing chemicals for which metabolism depends on the P450 enzyme system (Hoen et al., 2000; Boess et al., 2003). Human hepatocytes also tend to show a decrease in most P450 enzymes, but some levels of P450 gene expressions can be restored after a few days (Morel et al., 1990; LeCluyse, 2001). A mouse in vitro hepatocyte system might be an alternative to the rat and human systems, especially if the metabolic competence is preserved. In addition, because the complete sequence of the mouse genome is known (Waterston et al., 2002) and transgenic mouse models are widely available, primary mouse hepatocytes are usable for the purpose of performing mechanistic investigations of liver toxicity. However, information about stability of liver-specific functions in mice is scarce. It was the aim of the current study to investigate the robustness of a mouse hepatocyte-based in vitro model, especially with respect to the biotransformation functions (evaluated by assessing gene expression and enzyme activities), to gain more insight in its potential usefulness in toxicology.

### Materials and Methods

**Chemicals.** Dulbecco's modified Eagle's medium, fetal calf serum, penicillin/streptomycin, Hanks' calcium- and magnesium-free buffer, insulin, and TRIzol were obtained from Invitrogen (Breda, The Netherlands). Glucagon, hydrocortisone, collagenase type IV, dimethylsulfoxide (DMSO), and trypan blue were purchased from Sigma-Aldrich (Zwijndrecht, The Netherlands). Collagen Type I, rat tail was obtained from BD Biosciences Discovery Labware (Bedford, MA).

The RNeasy minikit was obtained from QIAGEN, Westburg B.V. (Leusden, The Netherlands). The 5× MEGAscript T7 Kit was obtained from Ambion (Austin, TX). The GeneChip Expression 3'-Amplification Two-Cycle cDNA Synthesis Kit and Reagents, the Hybridization, Wash, and Stain Kit, and the Mouse Genome 430 2.0 Arrays were purchased from Affymetrix (Santa Clara, CA). The P450 AL4 mixture was delivered by Advanced In Vitro Cell Technologies (Valencia, Spain).

**Animals.** Permission for animal studies was obtained from the Animal Ethical Committee. Adult male C57BL/6 mice, weighing 20 to 25 g, were obtained from Charles River GmbH (Sulzfeld, Germany). This mouse strain was chosen because it is frequently used in toxicological and pharmacological investigations, and it is a common background for transgenic mouse strains. The animals were housed in Macrolon (Technilab-BMI, Someren, The Netherlands) cages with sawdust bedding at 22°C and 50 to 60% humidity. The light cycle was 12-h light/12-h dark. Feed and tap water were available ad libitum.

**Isolation of Hepatocytes.** Hepatocytes were isolated from adult male C57BL/6 mice by a two-step collagenase perfusion method according to Seglen (1976) and Casciano (2000) with modifications. The liver was perfused after cannulation of the abdominal inferior vena cava. The hepatic portal vein was cut through, and the thoracic inferior vena cava was occluded with forceps. The organ was washed with Hanks' calcium- and magnesium-free buffer for 3 min. After the liver had been freed of blood, the calcium-free buffer was replaced by a collagenase buffer (0.5 mg/ml) for 7 to 10 min. A perfusion rate of 7 ml/min and a temperature around 39°C was maintained for both perfusates during the entire procedure. After the perfusion had been terminated, the liver was rapidly excised from the body cavity and transferred to a sterile Petri dish. The gall bladder and remnants of the diaphragm were removed, and cells were released by disrupting the liver capsule mechanically and by shaking the cells into attachment medium. The cells were separated from undigested tissue with a sterile 50-μm mesh nylon filter. After washing by low-speed centrifugation at 50g for 3 min at 4°C several times, cell viability and yield were determined by trypan blue exclusion.

**Cell Culture.** Cells from three independent biological replicates with viability >85% were used and cultured on collagen gel-precoated six-well plates at a density of  $6.5 \times 10^5$  cells/ml (Koebe et al., 1994). Cells were allowed to attach for 2 to 4 h at 37°C in a humidified chamber with 95:5% air/CO<sub>2</sub> in Dulbecco's modified Eagle's medium supplemented with 10% fetal calf serum, 0.5 U/ml insulin, 7 ng/ml glucagon, and 2% penicillin/streptomycin (5000

U/ml penicillin; 5000 μm/ml streptomycin). After attachment, medium containing unattached cells and debris was removed by washing, and the cultures were overlaid with a second collagen layer to form a collagen-collagen sandwich culture. Primary cultures of mouse hepatocytes were cultured at 37°C in a humidified chamber with 95:5% air/CO<sub>2</sub> in serum-free culture medium supplemented with 0.5 U/ml insulin, 7 ng/ml glucagon, 7.5 μg/ml hydrocortisone, and 2% penicillin/streptomycin (5000 U/ml penicillin; 5000 μm/ml streptomycin). Cells from three independent biological experiments (each from a different animal) were harvested at 0, 42, and 90 h after isolation for gene expression analysis.

**RNA Isolation.** Total RNA was isolated from cultured mouse hepatocytes using TRIzol reagent with the RNeasy kit according to the manufacturer's protocol. Total cellular levels were measured on a spectrophotometer, and the quality of each RNA preparation was determined with a bioanalyzer (Agilent Technologies, Amstelveen, The Netherlands). Extracted RNA was stored at -80°C.

**Target Preparation and Microarray Hybridization.** cRNA targets were prepared according to the Affymetrix protocol. The targets were hybridized according to the manufacturer's recommended procedures on high-density oligonucleotide GeneChips (Affymetrix Mouse Genome 430 2.0 GeneChip arrays). The GeneChips were washed and stained using a fluidics station by Affymetrix and scanned in an Affymetrix GeneArray scanner.

Nine targets (three time points from three animals) were prepared, and each preparation was analyzed using one GeneChip array. Normalization quality controls, including scaling factors, average intensities, present calls, background intensities, noise, and raw Q values all were within acceptable limits. Hybridization controls BioB, BioC, BioD, and CreX were called present on all the chips and yielded the expected increases in intensities.

**Data Analysis.** Nine datasets were obtained from this experiment. The raw data were imported into ArrayTrack (Tong et al., 2003) and converted and normalized by Robust Multi-array Average (integrated into ArrayTrack) (Irizarry et al., 2003). Spots of poor quality and control spots were omitted. Subsequently, the remaining probe sets (25,084) were logarithm (base 2)-transformed. Processed and normalized gene expression values were evaluated for identifying differentially expressed probe sets with the multiclass significance analysis of microarrays (SAM, integrated into ArrayTrack;  $p < 0.05$ ) (Shi et al., 2006). Multiclass SAM resulted in 6560 remaining differentially expressed probe sets, and these were used for principal component analysis (PCA).

**Pathway Analysis.** Gene expressions at 42 versus 0 h, 90 versus 0 h, and 90 versus 42 h were uploaded in T-profiler (<http://www.t-profiler.org>; Boersma et al., 2005) to identify transcriptional regulation of biochemical pathways and biological processes that included genes sets from Gene Ontology terms, curated gene sets, motifs, and Kyoto Encyclopedia for Genes and Genomes pathways. T-profiler uses the *t* test to score the difference between the mean expression level of predefined groups of genes and that of all the other genes without any statistical preselection of modulated genes (Boersma et al., 2005). Significance was determined by generating an E value, a Bonferroni-corrected *p* value. Pathways and processes were determined significant if E values were less than 0.05, and were thereupon used for hierarchical clustering and gene grouping in GenePattern (<http://www.broad.mit.edu/cancer/software/genepattern/>). To increase specificity of the transcriptomics changes, the 6560 differentially expressed probe sets, determined by multiclass SAM, were uploaded into MetaCore (GeneGo, San Diego, CA) for identifying the involvement of differentially expressed genes in specific cellular pathways by overrepresentation analyses compared with the total amount of genes involved in the particular pathways.

**Analysis of P450 Enzyme Activities.** P450 activity assays at 0, 42, and 90 h were conducted by the direct incubation of three independent sandwich-cultured mouse hepatocyte preparations with a mixture of P450 substrates (Lahoz et al., 2008). Substrate mixture stock was prepared in DMSO and conveniently diluted in incubation media to obtain the optimal assay concentrations (Table 1). The final concentration of DMSO during incubation was 0.5% (v/v). After 2 h of incubation, enzymatic reactions were stopped by aspirating the incubation medium. Samples were subsequently centrifuged at 2500 rpm for 5 min. The supernatants were transferred to clean tubes and frozen at -80°C until analysis. Sample analysis and mixture preparation was carried out in the Unidad Mixta Hospital La Fe-Advancell (Valencia, Spain).

TABLE 1  
Substrates in the P450 AL mixture for measurement of P450 activities

P450	Substrate	Concentration $\mu M$	Metabolite
1A2	Phenacetin	10	Acetaminophen
2A4	Coumarin	5	7-HO-Coumarin
2B9/10/13/19/23	Bupropion	10	HO-Bupropion
2C37	Mephenytoin	50	4'-HO-Mephenytoin
2C39-40/44/50/54-55/65-70	Diclofenac	10	4'-HO-Diclofenac
2D9-13/22/26/34/40	Bufuralol	10	Hydroxybufuralol
2E1	Chlorzoxazone	50	6-HO-Chlorzoxazone
3A11	Midazolam	5	1'-HO-Midazolam

Samples were subjected to hydrolysis before analysis by incubation with  $\beta$ -glucuronidase and arylsulfatase for 2 h at 37°C. One volume of ice-cold acetonitrile was added, and centrifuged samples were transferred to clean vials for further analysis. Metabolites formed and released into the culture medium were quantified by high-performance liquid chromatography/tandem mass spectrometry (Lahoz et al., 2007). Enzymatic activities were expressed as picomole of metabolite formed per incubation time per number of cells. For statistical analysis, the *t* test was used with a significance level of 5%.

## Results

**Principal Component Analysis of All Genes.** A PCA on the 6560 differentially expressed probe sets, identified with the multiclass SAM analysis ( $p < 0.05$ ), was used to visualize interindividual and time-dependent differences in gene expression profiles of primary mouse hepatocytes cultured in sandwich configuration (Fig. 1). PCA shows a clear discrimination between the three time groups, and only limited differences between replicates of the 42- and 90-h groups are observed, whereas more variance in gene expression is seen between the 0-h replicates.

**Pathway Analysis.** T-profiler was used for identifying gene groups that were differentially expressed. The gene groups significantly altered in hepatocytes from at least two experiments and modified in the same direction in liver cells from all the experiments were selected for each period. Three hundred eighty-four gene sets were identified as being modulated. Hierarchical clustering was used to generate more comprehensible groups of these genes (Fig. 2; Supplemental Data). Most gene sets were either only down-regulated or only up-regulated. A small amount of these gene sets seemed to be up-regulated during the first 42 h and down-regulated from 42 to 90 h.

An additional analysis of affected cellular pathways was conducted using MetaCore, which showed similar results compared with the results from T-profiler (data not shown). Most prominent pathways significantly altered in MetaCore were involved in several catabolic, metabolic, and biosynthetic processes, immune responses, and cellular, nuclear, and macromolecule-related processes.

**Biotransformation Enzymes.** Gene expression and activity of phase I and phase II drug-metabolizing enzymes were quantified to assess the metabolic capacity of the primary mouse hepatocyte system (see Tables 2 and 3). For the majority of the P450 genes, the gene expression level decreased during culturing. However, for a small group of P450s, the expression levels remained relatively constant, whereas for an even smaller group, gene expression levels increased in all the periods.

For the phase II genes, large variations were seen in their expression levels between the different time points, even within groups of phase II genes belonging to the same family. Most phase II enzyme groups were down-regulated or hardly changed over time. Only genes involved in glutathione metabolism and *N*-acetyltransferases were generally up-regulated.

P450 enzyme activities of eight selected enzymes were measured in




Fig. 1. Principal component analysis using the data of all the differentially expressed genes from primary mouse hepatocytes at 0, 42, and 90 h cultivation.

each sample. These activities are presented as  $\log_2$  mean  $\pm$  S.D. values in Fig. 3. Enzyme activities for Cyp2a4 and Cyp2a1 were not detected. For the majority of the studied P450 enzymes, a significant decrease in enzyme activity was measured in both periods. Only one group of enzymes (Cyp2B9-10/13/19/23) showed an increase in activity and expression levels. After 90 h in culture, all the P450 enzymes remained active in the primary hepatocytes.

These P450 enzyme activities and their gene expression are presented in Fig. 3. In all the cases gene expression and enzyme activity changed in the same directions, and mostly in the same order.

## Discussion

As human and rat sandwich-cultured hepatocytes have their disadvantages because of the limited availability of donor material and/or a rapid decline in P450 enzyme activities, primary mouse hepatocytes may be an option to overcome this shortcoming (Hoen et al., 2000; Schaeffner et al., 2005). Moreover, the growing availability of transgenic models can be a reason to use mouse primary hepatocytes. As information on the stability of liver-specific functions in mouse hepatocytes is scarce, the present study investigated the stability of a mouse hepatocyte-based in vitro model, especially with respect to the






FIG. 2. Hierarchical clustering of gene sets significantly modified in primary mouse hepatocytes in the time periods from 0 to 42 h, from 42 to 90 h, and from 0 to 90 h. Clustering is based on the *t* values, as provided by T-profiler. Yellow and blue mean up- and down-regulation, respectively.

biotransformation functions (gene expression and enzyme activities), to gain more insight in its potential usefulness in toxicology.

PCA was used to elucidate changes in gene expression profiles between donor animals and between cultivation times. This revealed that profiles are very different between the three time points. These time-dependent changes can be attributed to the adaptation of the primary hepatocytes to their in vitro environment. Noteworthy, the hepatocytes at time point 0 h show larger interanimal differences compared with those at later time points. This observation might be caused by initial stress induced by the enzymatic separation and isolation of hepatocytes (Pillar and Seitz, 1999). This interanimal variability is less clear in the later time points, indicating that the stress responses are fading.

Pathway analysis, used to identify biochemical pathways and biological processes that were differentially expressed, showed again that differences exist between the two cultivation periods. Some gene sets were affected in both periods, whereas others were affected in only the first 42 h or only in the 48 h thereafter. During only the first 42 h in culture, gene sets in mitochondrial processes, hormonal processes, and chemotaxis were down-regulated, whereas those in protein-related processes are up-regulated during only this period. Pathways mostly affected after the first 42 h are down-regulation of membrane-related processes and up-regulation of transcription and replication and cell

TABLE 2

Expression of P450 genes in primary mouse hepatocytes at 42 and 90 h as percentages of the expression at 0 h

Gene Name	42 h	90 h
Cyp17a1*	2659	107
Cyp1a1	119	117
Cyp1a2*	29	7
Cyp1b1	109	104
Cyp20a1	155	213
Cyp21a1	114	108
Cyp26a1*	24	10
Cyp27a1*	66	56
Cyp2a12*	65	128
Cyp2a4/5	42	71
Cyp2b10	1546	1460
Cyp2b13	102	95
Cyp2b19	90	86
Cyp2b9	126	110
Cyp2c29*	32	13
Cyp2c37	24	1
Cyp2c37/50/54*	19	1
Cyp2c38	54	65
Cyp2c44*	27	2
Cyp2c55*	50	16
Cyp2c70*	75	39
Cyp2d10*	88	59
Cyp2d13*	40	20
Cyp2d22*	74	33
Cyp2d26*	69	56
Cyp2d9*	59	22
Cyp2e1*	56	3
Cyp2f2*	22	4
Cyp2g1*	40	37
Cyp2j5*	10	6
Cyp2j6*	99	83
Cyp2r1	103	76
Cyp2s1*	266	83
Cyp2u1	76	80
Cyp39a1*	197	421
Cyp3a11*	77	76
Cyp3a13*	203	175
Cyp3a25*	21	18
Cyp3a41*	66	65
Cyp3a44*	82	77
Cyp46a1*	83	67
Cyp4a10*	32	38
Cyp4a12*	16	1
Cyp4a14*	19	7
Cyp4b1	202	100
Cyp4f13	70	58
Cyp4f14*	19	5
Cyp4f15*	26	9
Cyp4f16*	381	341
Cyp4v3*	41	22
Cyp51	115	108
Cyp7a1	76	42
Cyp7b1*	17	12
Cyp8b1*	109	7
	69	57

\* Significant ( $p < 0.05$ ) in multiclass SAM.

cycle-related processes. Gene sets that are continuously down-regulated are on humoral and cell-mediated immunity, P450-related responses, and liver-specific genes. A continuous up-regulation is observed in nuclear and nucleic acid-related metabolism, cellular organization, and in parts of cell cycle-related processes. The overall picture based on these pathway analyses is that liver-specific functions are continuously decreasing and that gross cellular organizations are continuously up-regulated. Initial effects are more at the level of protein synthesis, external signaling pathways, and energy production, whereas later transcription, translation, membranes, and cell cycle-related gene sets are affected.

Microscopic observations showed that in this 42-h period, cells

TABLE 3

Expression of phase II genes in primary mouse hepatocytes at 42 and 90 h as percentages of the expression at 0 h

Gene Name	42 h	90 h
Aldo-keto reductases		
Akr1a4	92	101
Akr1b3*	208	321
Akr1b7*	267	285
Akr1b8	131	127
Akr1c12	96	85
Akr1c13	103	107
Akr1c14*	29	47
Akr1c19	125	84
Akr1c20*	27	22
Akr1c21*	86	71
Akr1c6*	4	0
Akr1d1*	20	13
Akr1e1	112	103
Akr7a5	93	83
Sulfotransferases		
Chst10	81	72
Chst11	92	81
Chst4	83	80
Chst9	98	95
D4st1*	151	239
Gal3st1	175	128
Gal3st4	96	81
Hs1bp3*	120	205
Hs2st1*	167	229
Hs3st1	93	88
Hs3st2	88	75
Hs3st3b1*	88	76
Hs3st6	83	73
Hs6st1*	86	59
Hs6st2	94	89
Ndst1	99	100
Ndst2	165	113
Ndst3	94	92
Ndst4	93	84
Sult1a1	110	119
Sult1b1*	67	18
Sult1c2	76	50
Sult1d1*	457	86
Sult1e1*	1231	397
Sult5a1*	67	60
Epoxide hydrolases		
Ephx1	140	133
Ephx2*	47	35
Flavin-containing monooxygenase		
Fmo1*	31	19
Fmo2	102	96
Fmo3	99	84
Fmo4	133	108
Fmo5	369	96
Glutathione reductases		
Gsr	146	129
Gss*	153	329
Glutathione peroxidases		
Gpx1*	70	47
Gpx4*	139	161
Gpx6*	58	44
Gpx7*	113	222
Glutathione transferases		
Gsta1/2*	526	409
Gsta2*	194	142
Gsta3*	39	42
Gsta4*	82	47
Gstm1	113	127
Gstm2*	364	383
Gstm3*	378	593
Gstm4	127	125
Gstm5*	197	205
Gstm6	110	85
Gstm7*	64	51
Gsto1	117	97
Gsto2	115	115
Gstp1	98	107
Gstt1*	49	38

TABLE 3—Continued.

Gene Name	42 h	90 h
Gstt2	78	105
Gstt3	69	55
Gstz1*	94	92
Mgst1*	63	60
Mgst2*	94	76
Mgst3	141	150
N-Acetyltransferases		
Gcnt1	110	107
Gcnt2	76	63
Nat10*	153	239
Nat11	114	174
Nat12	145	196
Nat2*	179	122
Nat5	240	332
Nat6	82	114
Nat9	134	135
Glucuronosyl transferases		
Ugt1a1/2/5/6a/6b/7c/9/10*	59	38
Ugt2a3*	30	4
Ugt2b1*	13	1
Ugt2b34	127	112
Ugt2b35	99	147
Ugt2b36*	42	59
Ugt2b37*	40	28
Ugt2b38*	11	5
Ugt2b5*	35	39
Ugt3a1*	20	6
Ugt3a2*	26	4
	96	92

\* Significant ( $p < 0.05$ ) in multiclass SAM.

were changing their periphery, cellular components, and structures, and forming aggregates (data not shown). These morphological changes at the microscopic level confirm previously described studies on human hepatocytes and on the histology of rat and mouse hepatocytes. The morphological changes include attachment to the collagen layer, the remodeling of sandwich-cultured hepatocytes to a polygonal shape, the appearance of a clear, less granular cytoplasm, and the aggregation of the cells, including the formation of gap junctions for intercellular communication (Klaunig et al., 1981a,b; Tuschl and Mueller, 2006; Lu et al., 2007). Membrane and mitochondrial changes have also been shown before in freshly isolated hepatocytes and hepatocytes cultured for 3 days (Berry and Simpson, 1962; Ikeda et al., 1992). It is unfortunate that all of these morphological changes were poorly represented by changes at the level of pathways as analyzed by T-profiler or MetaCore. These morphological changes were perhaps too specific, and the annotation of genes with respect to that is too limited. In human and rat hepatocytes cultured in a sandwich configuration, others have found genes and pathways involved in cell structure to be highly expressed after 24 h in culture (Ben-Ze'ev et al., 1988; Schuetz et al., 1988; Liu et al., 1991; Waring et al., 2003).

To investigate the metabolic competence of sandwich-cultured mouse hepatocytes in more detail, gene expression profiles from phase I and phase II biotransformation enzymes were evaluated during the cultivation period. In general, the expression values of all the phase I biotransformation enzymes decreased over the whole cultivation period. During the first 42 h, these phase I enzyme gene expression values decreased by 31% (approximately 69% remaining) from the original values at 0 h. After 90 h in culture, the hepatocytes exhibit clearly lower levels (approximately 57%) of P450s compared with the original values at 0 h.

It was described before that many cultures of mammalian hepatocytes exhibit low levels of P450 but still possess the ability to respond to P450 inducers (Cervenkova et al., 2001). The expression of Cyp1a1

## Cyp450 activity vs expression




FIG. 3. P450 enzyme activities and gene expression in primary mouse hepatocytes after 0, 42, and 90 h in culture. Values are represented as percentages from activities and gene expression 0 h and as mean  $\pm$  S.D. of three animals for each time point. \*, significant ( $p < 0.05$ ) compared with 0 h; #, significant ( $p < 0.05$ ) compared with 0 and 42 h.

and Cyp1a2 during cultivation has been studied in detail by Northern blotting (Tamaki et al., 2005). In primary mouse hepatocytes cultured in dishes without collagen, Cyp1a1 expression increased during cultivation. Cyp1a2 was constitutively expressed in these cells, but insulin suppressed this expression slightly. Our results show a steady level in Cyp1a1 gene expression and a down-regulation in Cyp1a2 gene expression. In human hepatocytes, most P450s also showed a decrease in expression levels with the exception of a few P450s that had elevated levels of expression (Richert et al., 2006). Cyp1A2 seemed to be more stable compared with the mouse model during cultivation, whereas Cyp2E1 and Cyp3A4, which is the homolog for Cyp3a11 in mice (Nelson et al., 2004), were shown to decrease in time, comparable with our results (George et al., 1997). Most P450s in conventional sandwich-cultured rat hepatocytes seemed to decrease in a time-dependent manner (Kienhuis et al., 2007). Even within the short period of 24 h after isolation, the expression levels of many P450s decreased tremendously in rat hepatocytes (Boess et al., 2003), whereas in our study gene expression levels showed a slower decrease over time, indicating that mouse hepatocytes remain more stable with respect to P450 gene expression compared with rat hepatocytes but not with human hepatocytes.

Because several rat studies suggest that phase II enzyme activities are better preserved in culture than phase I enzyme activities, we decided to only measure enzyme activities of the most important phase I enzymes involved in xenobiotic biotransformation to strengthen our gene expression results (Rogiers and Vercruysse, 1993; Kern et al., 1997). These results show a general decrease in P450 enzyme activity, with only one exception. Activity levels decreased the first 42 h to 59% from the original activities. After 90 h, activity levels decreased to 34% compared with 0 h. Compared with our results, P450 enzyme activities in rat hepatocytes decrease very rapidly. Already after 6 h, less than 50% of the initial P450 enzyme activity was left in rat hepatocytes (Lopez-Garcia, 1998; Lopez-Garcia and Sanz-Gonzalez, 2000). However, these rat hepatocytes were cultured on a collagen monolayer, which may affect the stability of the cells.

Phase II biotransformation enzymes show a less clear pattern. Enzymes were either up- or down-regulated or remain unchanged. Even within a group of enzymes, differences in gene expression can be seen. In general, the expression levels decrease to 96 and 92% compared with levels at 0 h after 42 and 90 h, respectively. However, in rat hepatocytes, phase II enzymes in general showed a decrease in gene expression level after 72 h compared with gene expression in liver (Kienhuis et al., 2007). For rat hepatocytes, it is known that the

phase II enzymes are better preserved in culture than phase I enzymes (Rogiers and Vercruysse, 1993; Kern et al., 1997). Our data confirm these findings, thus showing that mouse hepatocytes are also stable in expression of genes encoding phase II enzymes. An overall comparison of the gene expression levels from phase I and phase II enzymes from rat and mouse hepatocytes showed that mouse hepatocytes exhibit a more stable gene expression pattern than primary rat hepatocytes.

### Conclusion

In summary, our findings show that the sandwich-cultured primary mouse hepatocyte in vitro model has little or no interanimal variations, showing its reproducibility. The system is still metabolic-competent even after 90 h of culturing, despite that phase I enzyme gene expression and activity generally decrease. In rat hepatocytes, this decrease in phase I and phase II gene expression appears to be faster. Human hepatocytes, on the other hand, also show a decrease in these phase I enzymes but are able to restore the gene expression levels of some P450s. Taken together, our results indicate that the primary mouse hepatocyte system is relatively stable, might maintain its metabolic competence longer than rat hepatocytes, and is roughly similar to that of human hepatocytes. Based on our gene expression data, we recommend applying a 42-h recovery period before using the mouse hepatocytes.

**Acknowledgments.** We thank Sandra Montero at Unidad Mixta Hospital La Fe-Advancell for technical assistance in P450 activity measurements.

### References

- Ben-Ze'ev A, Robinson GS, Bucher NL, and Farmer SR (1988) Cell-cell and cell-matrix interactions differentially regulate the expression of hepatic and cytoskeletal genes in primary cultures of rat hepatocytes. *Proc Natl Acad Sci U S A* **85**:2161–2165.
- Berry MN, Grivell AR, Grivell MB, and Phillips JW (1997) Isolated hepatocytes—past, present and future. *Cell Biol Toxicol* **13**:223–233.
- Berry MN and Simpson FO (1962) Fine structure of cells isolated from adult mouse liver. *Cell Biol* **15**:9–17.
- Blaauboer BJ, Boobis AR, Castell JV, Coecke S, Groothuis GMM, Guillouzo A, Hall TJ, Haworth GM, Lorenzon G, Miltenburger HG, et al. (1994) The practical applicability of hepatocyte cultures in routine testing. The report and recommendations of ECVAM Workshop 1. *ATLA* **22**:231–241.
- Boess F, Kamber M, Romer S, Gasser R, Muller D, Albertini S, and Suter L (2003) Gene expression in two hepatic cell lines, cultured primary hepatocytes, and liver slices compared with the in vivo liver gene expression in rats: possible implications for toxicogenomics use of in vitro systems. *Toxicol Sci* **73**:386–402.
- Boersma A, Foat BC, Vis D, Klis F, and Bussemaker HJ (2005) T-profiler: scoring the activity of predefined groups of genes using gene expression data. *Nucleic Acids Res* **33**:W592–W595.
- Casciano DA (2000) Development and utilization of primary hepatocyte culture systems to evaluate metabolism, DNA binding, and DNA repair of xenobiotics. *Drug Metab Rev* **32**:1–13.
- Cervenkova K, Belejova M, Vesely J, Chmela Z, Rypka M, Ulrichova J, Modriansky M, and

- Maurel P (2001) Cell suspensions, cell cultures, and tissue slices—important metabolic in vitro systems. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* **145**:57–60.
- Davila JC, Rodriguez RJ, Melchert RB, and Acosta D Jr (1998) Predictive value of in vitro model systems in toxicology. *Annu Rev Pharmacol Toxicol* **38**:63–96.
- George J, Goodwin B, Liddle C, Tapner M, and Farrell GC (1997) Time-dependent expression of cytochrome P450 genes in primary cultures of well-differentiated human hepatocytes. *J Lab Clin Med* **129**:638–648.
- Hoen PA, Commandeur JN, Vermeulen NP, Van Berkel TJ, and Bijsterbosch MK (2000) Selective induction of cytochrome P450 3A1 by dexamethasone in cultured rat hepatocytes: analysis with a novel reverse transcriptase-polymerase chain reaction assay section sign. *Biochem Pharmacol* **60**:1509–1518.
- Ikedo T, Ishigamia A, Anzai K, and Goto S (1992) Change with donor age in the degradation rate of endogenous proteins of mouse hepatocytes in primary culture. *Arch Gerontol Geriatr* **15**:181–188.
- Irizarry RA, Hobbs B, Collin F, Beazer-Barclay YD, Antonellis KJ, Scherf U, and Speed TP (2003) Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics* **4**:249–264.
- Kern A, Bader A, Pichlmayr R, and Sewing KF (1997) Drug metabolism in hepatocyte sandwich cultures of rats and humans. *Biochem Pharmacol* **54**:761–772.
- Kienhuis AS, Wortelboer HM, Maas WJ, van Herwijnen M, Kleinjans JC, van Delft JH, and Stierum RH (2007) A sandwich-cultured rat hepatocyte system with increased metabolic competence evaluated by gene expression profiling. *Toxicol In Vitro* **21**:892–901.
- Klaunig JE, Goldblatt PJ, Hinton DE, Lipsky MM, Chacko J, and Trump BF (1981a) Mouse liver cell culture. I. Hepatocyte isolation. *In Vitro* **17**:913–925.
- Klaunig JE, Goldblatt PJ, Hinton DE, Lipsky MM, and Trump BF (1981b) Mouse liver cell culture. II. Primary culture. *In Vitro* **17**:926–934.
- Koebe HG, Wick M, Cramer U, Lange V, and Schildberg FW (1994) Collagen gel immobilisation provides a suitable cell matrix for long term human hepatocyte cultures in hybrid reactors. *Int J Artif Organs* **17**:95–106.
- Lahoz A, Donato MT, Castell JV, and Gómez-Lechón MJ (2008) Strategies to in vitro assessment of major human CYP enzyme activities by using liquid chromatography tandem mass spectrometry. *Curr Drug Metab* **9**:12–19.
- Lahoz A, Donato MT, Picazo L, Gómez-Lechón MJ, and Castell JV (2007) Determination of major human cytochrome P450s activities in 96-well plates using liquid chromatography tandem mass spectrometry. *Toxicol In Vitro* **21**:1247–1252.
- LeCluyse EL (2001) Human hepatocyte culture systems for the in vitro evaluation of cytochrome P450 expression and regulation. *Eur J Pharm Sci* **13**:343–368.
- LeCluyse EL, Ahlgren-Beckendorf JA, Carroll K, Parkinson A, and Johnson J (2000a) Regulation of glutathione S-transferase enzymes in primary cultures of rat hepatocytes maintained under various matrix configurations. *Toxicol In Vitro* **14**:101–115.
- LeCluyse EL, Fix JA, Audus KL, and Hochman JH (2000b) Regeneration and maintenance of bile canalicular networks in collagen-sandwiched hepatocytes. *Toxicol In Vitro* **14**:117–132.
- Liu JK, DiPersio CM, and Zaret KS (1991) Extracellular signals that regulate liver transcription factors during hepatic differentiation in vitro. *Mol Cell Biol* **11**:773–784.
- López-García MP (1998) Endogenous nitric oxide is responsible for the early loss of P450 in cultured rat hepatocytes. *FEBS Lett* **438**:145–149.
- López-García MP and Sanz-González SM (2000) Peroxynitrite generated from constitutive nitric oxide synthase mediates the early biochemical injury in short-term cultured hepatocytes. *FEBS Lett* **466**:187–191.
- Lu P, Prost S, Caldwell H, Tugwood JD, Betton GR, and Harrison DJ (2007) Microarray analysis of gene expression of mouse hepatocytes of different ploidy. *Mamm Genome* **18**:617–626.
- Morel F, Beaune PH, Ratanasavanh D, Flinois JP, Yang CS, Guengerich FP, and Guillouzo A (1990) Expression of cytochrome P-450 enzymes in cultured human hepatocytes. *Eur J Biochem* **191**:437–444.
- Nelson DR, Zeldin DC, Hoffman SM, Maltais LJ, Wain HM, and Nebert DW (2004) Comparison of cytochrome P450 (CYP) genes from the mouse and human genomes, including nomenclature recommendations for genes, pseudogenes and alternative-splice variants. *Pharmacogenetics* **14**:1–18.
- Pillar TM and Seitz HJ (1999) Oxidative stress response induced in rat primary hepatocyte monolayers by mechanical removal of adherent cells. *Cell Tissue Res* **295**:363–367.
- Richert L, Binda D, Hamilton G, Viollon-Abadie C, Alexandre E, Bigot-Lasserre D, Bars R, Coassolo P, and LeCluyse E (2002) Evaluation of the effect of culture configuration on morphology, survival time, antioxidant status and metabolic capacities of cultured rat hepatocytes. *Toxicol In Vitro* **16**:89–99.
- Richert L, Liguori MJ, Abadie C, Heyd B, Mantion G, Halkic N, and Waring JF (2006) Gene expression in human hepatocytes in suspension after isolation is similar to the liver of origin, is not affected by hepatocyte cold storage and cryopreservation, but is strongly changed after hepatocyte plating. *Drug Metab Dispos* **34**:870–879.
- Rogiers V and Verbruggen A (1993) Rat hepatocyte cultures and co-cultures in biotransformation studies of xenobiotics. *Toxicology* **82**:193–208.
- Schaeffner I, Petters J, Aurich H, Froberg P, and Christ B (2005) A microtiterplate-based screening assay to assess diverse effects on cytochrome P450 enzyme activities in primary rat hepatocytes by various compounds. *Assay Drug Dev Technol* **3**:27–38.
- Schuetz EG, Li D, Omiecinski CJ, Muller-Eberhard U, Kleinman HK, Elswick B, and Guzelian PS (1988) Regulation of gene expression in adult rat hepatocytes cultured on a basement membrane matrix. *J Cell Physiol* **134**:309–323.
- Seglen PO (1976) Preparation of isolated rat liver cells. *Methods Cell Biol* **13**:29–83.
- Shi L, Reid LH, Jones WD, Shippy R, Warrington JA, Baker SC, Collins PJ, de Longueville F, Kawasaki ES, Lee KY, et al. (2006) The MicroArray Quality Control (MAQC) project shows inter- and intraplatform reproducibility of gene expression measurements. *Nat Biotechnol* **24**:1151–1161.
- Tamaki H, Sakuma T, Uchida Y, Jaruchotikamol A, and Nemoto N (2005) Activation of CYP1A1 gene expression during primary culture of mouse hepatocytes. *Toxicology* **216**:224–231.
- Tong W, Cao X, Harris S, Sun H, Fang H, Fuscoe J, Harris A, Hong H, Xie Q, Perkins R, et al. (2003) ArrayTrack—supporting toxicogenomic research at the U.S. Food and Drug Administration National Center for Toxicological Research. *Environ Health Perspect* **111**:1819–1826.
- Tuschl G and Mueller SO (2006) Effects of cell culture conditions on primary rat hepatocytes—cell morphology and differential gene expression. *Toxicology* **218**:205–215.
- Waring JF, Ciurlionis R, Jolly RA, Heindel M, Gagne G, Fagerland JA, and Ulrich RG (2003) Isolated human hepatocytes in culture display markedly different gene expression patterns depending on attachment status. *Toxicol In Vitro* **17**:693–701.
- Waterston RH, Lindblad-Toh K, Birney E, Rogers J, Abril JF, Agarwal P, Agarwala R, Ainscough R, Alexandersson M, An P, et al. (2002) Initial sequencing and comparative analysis of the mouse genome. *Nature* **420**:520–562.

**Address correspondence to:** Joost H. M. van Delft, Department of Health Risk Analyses and Toxicology, Faculty of Health, Medicine, and Life Sciences, Maastricht University, Universiteitssingel 50, 6229 ER Maastricht, The Netherlands. E-mail: j.vandelft@grat.unimaas.nl