

# NIH Public Access

Author Manuscript

J Cardiovasc Electrophysiol. Author manuscript; available in PMC 2012 September 1.

Published in final edited form as:

*J Cardiovasc Electrophysiol*. 2011 September ; 22(9): 1057–1066. doi:10.1111/j. 1540-8167.2011.02053.x.

## Comparison of the Effects of a Transient Outward Potassium Channel Activator on Currents Recorded from Atrial and Ventricular Cardiomyocytes

Kirstine Calloe, PhD<sup>3</sup>, Eyal Nof, MD<sup>2</sup>, Thomas Jespersen, PhD<sup>3</sup>, José M Di Diego, MD<sup>1</sup>, Natalie Chlus<sup>1</sup>, Søren-Peter Olesen, MD, PhD<sup>3</sup>, Charles Antzelevitch, PhD<sup>1</sup>, and Jonathan M Cordeiro, PhD<sup>1</sup>

<sup>1</sup> Department of Experimental Cardiology, Masonic Medical Research Laboratory Utica, New York, USA

<sup>2</sup> Leviev Heart Center, Sheba Medical Center, Tel Hashomer, Ramat Gan, Israel

<sup>3</sup> Danish National Research Foundation Center for Cardiac Arrhythmias, University of Copenhagen, Copenhagen, Denmark

## Abstract

**Introduction**—NS5806 activates the transient outward potassium current ( $I_{to}$ ) in canine ventricular cells. We compared the effects of NS5806 on canine atrial versus ventricular tissues and myocytes.

**Methods and Results**—NS5806 (10  $\mu$ M) was evaluated in arterially-perfused canine right atrial and right ventricular wedge preparations. In ventricular wedges NS5806 (10  $\mu$ M) accentuated phase 1 in epicardium (Epi), with little effect in endocardium (Endo), resulting in augmented J-waves on the ECG. In contrast, application of NS5806 (10  $\mu$ M) to atrial preparations had no effect on phase 1 repolarization but significantly decreased upstroke velocity (dV/dt) and depressed excitability, consistent with sodium channel block. Current and voltage-clamp recordings were made in the absence and presence of NS5806 in (10  $\mu$ M) enzymatically dissociated atrial and ventricular myocytes. In ventricular myocytes, NS5806 increased I<sub>to</sub> magnitude by 80% and 16% in Epi and Endo, respectively (at +40 mV). In atrial myocytes, NS5806 increased peak I<sub>to</sub> by 25% and had no effect on the sustained current, I<sub>Kur</sub>. Under control conditions, I<sub>Na</sub> density in atrial myocytes was nearly double that in ventricular myocytes. NS5806 caused a shift in steady-state mid-inactivation (V<sub>1/2</sub>) from  $-73.9\pm0.27$  to  $-77.3\pm0.21$  mV in ventricular and from  $-82.6\pm0.12$  to  $-85.1\pm0.11$  mV in atrial cells, resulting in reduction of I<sub>Na</sub> in both cell types. Expression of mRNA encoding putative I<sub>Na</sub> and I<sub>to</sub> channel subunits was evaluated by qPCR.

**Conclusion**—NS5806 produces a prominent augmentation of  $I_{to}$  with little effect on  $I_{Na}$  in the ventricles, but a potent inhibition of  $I_{Na}$  with little augmentation of  $I_{to}$  in atria.

#### Keywords

Transient outward potassium current; sodium current; atria; ventricle; NS5806

AUTHOR CONTRIBUTIONS

Conception and design of study; JMC and KC. Collection, analysis and interpretation of data; JMC, KC, NC, EN, TJ, JMD. Drafting the article or revising it critically; JMC, KC, SPO, CA. All authors approved the final version of the manuscript. **Disclosures:** SPO is a consultant to NeuroSearch A/S.

Corresponding Author: Jonathan M Cordeiro, Ph.D., Masonic Medical Research Laboratory, 2150 Bleecker Street, Utica, NY 13501, U.S.A., Phone: (315) 735-2217 x132, Fax: (315) 735-5648, jcordeiro@mmrl.edu.

## INTRODUCTION

The action potential (AP) in cardiac muscle is an important physiological parameter: 1) it is responsible for electrical conduction, 2) it modulates the refractory period and 3) associated with each action potential is a contraction via the process of excitation-contraction coupling. Marked differences in the shape and duration of the cardiac action potential have been described in different regions of the mammalian heart, reflecting differences in the underlying ionic currents. For example, comparative studies of K<sup>+</sup> currents in atrial and ventricular tissue have shown that the inward rectifier K<sup>+</sup> current is large in ventricle versus atria<sup>1,2</sup> and the ultra rapid delayed rectifier K current, I<sub>Kur</sub>, is large in atria but small<sup>3</sup> or virtually absent in the ventricle<sup>4</sup>. These differences in the complement of K<sup>+</sup> currents between the 2 tissue types have been ascribed to variations in the molecular subunits<sup>5–7</sup>. Marked differences in the kinetics and properties of Na<sup>+</sup> current have also been described in the 2 tissue types<sup>8–10</sup>. Na<sup>+</sup> current in atrial tissue has a larger current density and more negative steady-state inactivation compared to ventricular tissue; these differences are thought to contribute to the atrial-selective anti-arrhythmic effects of many compounds.

In atrial and ventricular tissue from dog, the action potential exhibits a large, rapid phase 1 repolarization, suggesting the presence of a Ca<sup>2+</sup>-independent transient outward K<sup>+</sup> current (I<sub>to</sub>). I<sub>to</sub> is thought to be mediated by K<sub>v</sub>4.3<sup>11,12</sup> together with K<sup>+</sup> Channel Interacting Protein (KChIP2)<sup>13</sup>. The expression of KChIP2 is high in the epicardium (Epi) and midmyocardium (Mid), and low in the endocardium (Endo) which is believed to account for the transmural gradient of I<sub>to</sub><sup>14,15</sup> but other proteins, such as the dipeptidyl aminopeptidase-like protein (DPP6)<sup>15,16</sup> and KCNEs<sup>17,18</sup> may also associate with K<sub>v</sub>4.3 and affect current. It has been suggested that K<sub>v</sub>1.4 contributes to canie ventricular I<sub>to</sub> and although K<sub>v</sub>1.4 is uniformly expressed across the ventricular wall<sup>15,19</sup>, it may have a relatively larger contribution to Endo I<sub>to</sub>, where KChIP2 expression is lower, resulting in smaller K<sub>v</sub>4.3/KChIP2 current compared to the Epi- and Mid. The slow recovery from inactivation of Endo I<sub>to</sub> compared to Epi- and Mid<sup>15</sup> provide further support for a role of K<sub>v</sub>1.4. The molecular composition of the channels contributing to phase 1 repolarization in the canine atria is also debated. Besides I<sub>to</sub>, a slower inactivating, ultra rapid delayed rectifier current (I<sub>Kur</sub>) is present. It is believed that K<sub>v</sub>1.5 mediates canine I<sub>Kur</sub> similar to human atrial I<sub>Kur</sub><sup>6</sup>.

We have recently identified a compound, NS5806 that produces an increase in the size of current carried through  $K_v4.3/KChIP2$  channels expressed in both *Xenopus laevis* oocytes and CHO-K1 cells. Similarly, this compound activates  $I_{to}$  in canine ventricular Epi- and Mid cells, whereas minimal effects on Endo  $I_{to}$  were observed. In this study, we investigated the effect of NS5806 on canine atrial currents compared to effects on ventricular currents. Application of NS5806 resulted in about a 30% increase in the magnitude of  $I_{to}$  in atria, a 80% increase in Epi  $I_{to}$ , and no significant effect on Endo  $I_{to}$  and. Surprisingly, application of NS5806 caused a marked reduction in the magnitude of Na<sup>+</sup> current in atrial cells compared to ventricular cells. Preliminary results have been presented in abstract form<sup>20</sup>.

## METHODS

#### NS5806

NS5806 was synthesized at NeuroSearch A/S, Ballerup, Denmark by reaction of 6cyano-2,4-dibromoaniline with sodium azide to form the respective 2,4-dibromo-6tetrazolylaniline, which was condensed with 3,5-bis-trifluoromethyl-phenylisocyanate to provide the target 1-[2,4-dibromo-6-(1H-tetrazol-5-yl)-phenyl]-3-(3,5-bis-trifluoromethylphenyl)-urea (NS5806). NS5806 was dissolved in DMSO (20 mM stock).

#### **Experimental animals**

This investigation conforms to the Guide for Care and Use of Laboratory Animals published by the National Institutes of Health (NIH publication No 85–23, Revised 1996) and was approved by the Animal Care and Use Committee of the Masonic Medical Research Laboratory. Adult mongrel dogs of either sex were used to perform all experimental series. The animals were anticoagulated with heparin and anesthetized with pentobarbital (35mg/ kg, i.v.). Their hearts were rapidly removed and placed in cold (4°C) cardioplegic solution (in mM): NaCl 129, KCl 12, NaH<sub>2</sub>PO<sub>4</sub> 0.9, NaHCO<sub>3</sub> 20, CaCl<sub>2</sub> 1.8, MgSO<sub>4</sub> 0.5, glucose 5.5.

#### Isolated ventricular and atrial preparations

Experiments were performed using isolated arterially perfused canine right atrial (RA) preparations or right ventricular (RV) wedge preparations. The RV wedges  $(2 \times 1.5 \times 1 \text{ cm})$ were excised from the base of the right ventricular free wall. A descending branch of the right coronary artery (RCA) was cannulated to deliver the perfusate. The isolated atrial preparations were perfused through the ostium of the RCA. During the cannulation procedure, the preparations were perfused with cardioplegic solution (4° C Tyrode's containing 12 mM KCl) and arterial branches were ligated using silk ligatures. Subsequently, the preparations were placed in a tissue bath and perfused with Tyrode's solution of the following composition (mM): NaCl 129, KCl 4, NaH<sub>2</sub>PO<sub>4</sub> 0.9, NaHCO<sub>3</sub> 20, CaCl<sub>2</sub> 1.8, MgSO<sub>4</sub> 0.5, glucose 5.5, bubbled with 95% O <sub>2</sub> and 5% CO<sub>2</sub> ( $37\pm0.5^{\circ}$ C). The perfusate was delivered at a constant flow. Pacing stimuli was delivered to the Endo surface at 2x the diastolic excitability at a basic cycle length (BCL) of 1 s. Control recordings were obtained 1-1.5 hours after mounting; the compounds were applied for 30 min before recordings. A pseudo-ECG was recorded with 2 electrodes consisting of Ag/AgCl half-cells placed 1.0 to 1.2 cm from 2 opposite sides of the atrial preparation. In the RV ventricular wedge, the electrodes were position facing the Endo and Epi surfaces (Epi, positive pole). Intracellular recordings were obtained using floating glass microelectrodes (Endo, approx. 2-3 mm from the endocardial surface, Epi, approx 0-3 mm from the Epi surface, atria, from the pectinate muscle (PM) and the crista terminalis (CT)). ECG and AP signals were amplified and digitized and analyzed using Spike 2 for Windows (Cambridge Electronic Design [CED], Cambridge, UK).

#### Isolation of adult canine cardiomyocytes

Myocytes from the Epi and Endo region were prepared from canine ventricles or atria. Left ventricular or right atrial preparations were dissected as described above. The preparation was initially perfused with nominally  $Ca^{2+}$ -free solution (mM): NaCl 129, KCl 5.4, MgSO<sub>4</sub> 2.0, NaH<sub>2</sub>PO<sub>4</sub> 0.9, glucose 5.5, NaHCO<sub>3</sub> 20, bubbled with 95% O<sub>2</sub>/5% CO<sub>2</sub> containing 0.1% BSA for a period of about 5 minutes. The preparation was then subjected to enzyme digestion with the nominally  $Ca^{2+}$ -free solution supplemented with 0.5 mg/ml collagenase (Type II, Worthington), 0.1 mg/ml protease (Type XIV, Sigma) and 1 mg/ml BSA for 8–12 minutes. For isolation of ventricular cells, thin slices of tissue from the Epi and Endo were shaved from the wedge using a dermatome. For atrial cells, the pectinate muscle was isolated. The tissue was then placed in separate beakers minced and incubated in fresh buffer containing 0.5 mg/ml collagenase, 1 mg/ml BSA and agitated. The supernatant was filtered, centrifuged at 200 rpm for 2 minutes and the myocyte containing pellet was stored in 0.5 mM Ca<sup>2+</sup> HEPES buffer at room temperature.

#### Voltage Clamp Recordings

Cells were placed in a temperature controlled chamber (PDMI-2, Medical Systems Corp.) mounted on the stage of an inverted microscope (Nikon TE300). Voltage-clamp and current-

clamp recordings were made using a MultiClamp 700A amplifier and MultiClamp Commander (Axon Instruments). Patch pipettes were fabricated from borosilicate glass capillaries (1.5 mm O.D., Fisher Scientific, Pittsburgh, PA). The pipettes were pulled using a gravity puller (Model PP-830, Narishige, Tokyo, Japan) and the pipette resistance ranged from 1–3 M $\Omega$ . Initially the cardiomyocytes were superfused with a HEPES buffer of the following composition (mM): NaCl 126, KCl 5.4, MgCl<sub>2</sub> 1.0, CaCl<sub>2</sub> 2.0, HEPES 10, and glucose 11. pH adjusted to 7.4 with NaOH. The patch pipette solution had the following composition (mM): K-aspartate 90, KCl 30, glucose 5.5, MgCl<sub>2</sub> 1.0, EGTA 5, MgATP 5, HEPES 5, NaCl 10, pH=7.2 with KOH and APs were recorded in this solution. For Ito recordings the HEPES buffer was supplemented with 300 µM cadmium to block I<sub>Ca</sub> as previously described<sup>21</sup>. Sodium currents were recorded as previously described<sup>22</sup> in a low Na<sup>+</sup> extracellular solution (in mM): CaCl<sub>2</sub> 0.5, Glucose 10, MgCl<sub>2</sub> 1.5, Choline-Cl 120, NaCl 5, HEPES 10, KCl 4, Na acetate 2.8, CoCl<sub>2</sub> 1, BaCl<sub>2</sub> 0.1, pH=7.4. The pipette solution consisted of: MgCl<sub>2</sub> 1, NaCl 15, KCl 5, CsF 120, HEPES 10, EGTA 10, Na<sub>2</sub>ATP 4, and the pH was adjusted to 7.2. All experiments were performed at 37° C except for the sodium current recordings which were performed at room temperature. After a whole cell patch was established, cell capacitance was measured by applying -5 mV voltage steps. Electronic compensation of series resistance to 60-70% was applied to minimize voltage errors. All analog signals (cell current and voltage) were acquired at 10-50 kHz, filtered at 4-6 kHz, digitized with a Digidata 1322 converter (Axon Instruments) and stored using pClamp9

#### **Quantitative Real Time PCR**

software

mRNA extraction and real-time reverse transcriptase-polymerase chain reaction (RT-PCR) was performed as previously reported<sup>15</sup>. The pre-designed gene expression assay from Applied Biosystems was, in addition to the previously reported, cf02625015 (SCN5A). The primers and probes targeting *SCN1B*, *SCN2B*, *SCN3B*, and *SCN4B* were designed and synthesized by Applied Biosystems, following submission of sequences including intron-exon boundaries using Primer Express 3.0 software.

## **Statistical Analysis**

Data are presented as Mean $\pm$ SEM. Statistical comparisons were made using paired or unpaired Student's t-test or one way ANOVA followed by Student Newman-Keuls post test for multiple comparison. p < 0.05 was considered statistically significant (\*).

#### RESULTS

As an initial basis of comparison, we tested the effect of NS5806 in arterially perfused right atrial preparations (Figure 1A). Application of 10  $\mu$ M NS5806 had no significant effect on phase 1 repolarization (from 5.1±2.1% in control to 6.3±3.7% of phase 2 amplitude, n=4). Action potential duration (APD) was virtually unchanged following 10  $\mu$ M NS5806 (APD<sub>90</sub>=223±9.7 ms before and 247±13.8 ms after application of NS5806, n=4, p=N.S.) and the effective refractory period was158±11 ms in control and 183±10 ms in presence of NS5806 (n=4, p=N.S.). At 15  $\mu$ M NS5806, the atrial action potential showed a depressed phase 0 depolarization and in 2 out of 4 preparations, the depressed phase 0 depolarization deteriorated into complete loss of excitability suggesting block of sodium channels. Similar effects were found on action potentials recorded from the crista terminalis (data not shown).

In right ventricular preparations, the Epi APs have a prominent phase 1 repolarization and this was further increased by NS5806 (10  $\mu$ M). In contrast, Endo APs have a small phase 1 repolarization and the APs were unaffected by 10  $\mu$ M NS5806. The increase in the Epi notch size was accompanied by an accentuation of the J-wave on the ECG (Figure 1B). At 15  $\mu$ M

To further compare the effect of NS5806 on canine atrial and ventricular cells, we characterized the effect of NS5806 on APs recorded from isolated cardiomyocytes. Figure 1C shows APs and corresponding upstroke velocity (dV/dt) in the absence and presence of NS5806. Application of 10  $\mu$ M NS5806 to isolated right atrial cells (Figure 1C, left panel) dramatically reduced upstroke velocity (dV/dt) by 71±7% (n=7) resulting in a decrease in phase 0 amplitude and in APD<sub>90</sub> (Table 1). For Epi cells, application of NS5806 resulted in a dramatic increase in phase 1 repolarization leading to loss of the AP dome, which was accompanied by a significant abbreviation of the APD<sub>90</sub>. The dV/dt was decreased by 9.8±4.8% (n=5) and phase 0 amplitude unaffected (Figure 1C, middle panel). NS5806 did not have any significant effect on the waveform of Endo APs and dV/dt was reduced by 4.9±2.8% (n=4, p=N.S.) (Figure 1C, right panel). The effect on the different electrophysiological parameters is summarized in Table 1.

To address the differences in the effect of NS5806 (10  $\mu$ M) on phase 1 repolarization in atrial and ventricular cells, the density of I<sub>to</sub> was measured in isolated canine atrial cells and ventricular Endo and Epi cells. I<sub>to</sub> was recorded using whole-cell patch-clamp techniques and Cd<sup>2+</sup> (300  $\mu$ M) was added to the extracellular solution to block I<sub>CaL</sub>. Following a brief step to -50 mV to discharge sodium channels, voltage steps from -40 to +50 mV were applied and elicited fast activating and rapidly inactivating currents as well as a sustained pedestal current component (Figure 2A). Application of NS5806 significantly increased the peak current density from 15.5±1.6 to 18.9±2.3 pA/pF (at +50 mV, n=14) whereas NS5806 did not affect the sustained current component (from 3.8±0.3 to 3.7±0.3 pA/pF) (Figure 2B). Mono-exponential equations were fit to the current traces and the time constants of decay ( $\tau$ ) following application of NS5806 were significantly slower (from  $\tau$ =20.0±0.7 to 26.5±1.5 ms at +50 mV) (Figure 2C). For comparison, currents recorded from ventricular Epi and Endo cells using identical experimental conditions are shown in Figure 2D-I.

The steady-state inactivation of atrial I<sub>to</sub> was evaluated using a prepulse-test pulse voltageclamp protocol in the presence of Cd<sup>2+</sup> and the effect of 10  $\mu$ M NS5806 was evaluated (Figure 3A and B). The peak current following a 2 s prepulse was normalized to the maximum current and plotted as a function of the prepulse voltage to obtain the availability of the channels. A Boltzmann function was fit to the data (Figure 3C). In the absence of drug, the mid-inactivation voltage was V<sub>1/2</sub>=-33.8±0.46 mV and the slope factor k= -7.15±0.40. Application of NS5806 caused a significant negative shift in V<sub>1/2</sub> to -41.1±0.50 mV and k=-7.35±0.42, (n=8). By comparison, the mid-inactivation voltage of I<sub>to</sub> was -46.0±1.0 mV, k=-4.75±0.44 for LV Epi cells and -52.9±1.1, k=-8.45±1.34 for LV Endo cells. Application of NS5806 (10  $\mu$ M) significantly shifted V<sub>1/2</sub> to -52.0±0.9 mV, k= -4.18±0.37 for Epi and -56.3±0.8 mV, k=-7.73±1.07 for Endo cells.

We next tested if recovery from inactivation of atrial  $I_{to}$  was affected by 10  $\mu$ M NS5806. Currents were elicited by a double pulse protocol, with increasing interpulse intervals. In absence of NS5806, reactivation of  $I_{to}$  showed a fast and a slow phase, with  $\tau_{fast}$  of 55±1.97 ms and a  $\tau_{slow}$  of 2368±327 ms. In the presence of NS5806, the reactivation of atrial  $I_{to}$  conformed to a single exponential equation and was markedly faster with  $\tau$ =69.3±1.0 ms (n=5). The results are summarized in Figure 3D.

In contrast to Epi cells, atrial cells have a prominent sustained current,  $I_{Kur}$ . To address the effect of NS5806 on the sustained current, a two-pulse protocol was used to separate  $I_{Kur}$ 

and I<sub>to</sub>. From a holding of -80 mV, I<sub>to</sub> was activated and inactivated by a step to 0 mV for 80 ms. The second pulse was preceded by a 5 ms step to -80 mV and I<sub>Kur</sub> was elicited by a series of voltages ranging from -30 to 40 mV in absence and presence of 10  $\mu$ M NS5806<sup>23</sup>. Application of NS5806 did not alter peak or sustained I<sub>Kur</sub> (data not shown).

Application of NS5806 to atrial cells resulted in a decrease in dV/dt (Figure 1C and Table 1) suggesting an inhibition of atrial sodium channels. We next recorded  $I_{Na}$  from cells isolated from canine atrial or ventricular Endo. Endo cells were chosen to minimize the impact of  $I_{to}$ . Representative  $I_{Na}$  recordings are shown in Figure 4. The current-voltage relationship shows a two-fold higher  $I_{Na}$  density in atrial cells compared to Endo cells. Steady-state gating parameters were addressed using a standard prepulse-testpulse voltage protocol and mid-inactivation potential ( $V_{l_2}$ ) of  $-82.6\pm0.12$  mV (n=8) for atrial and  $-73.9\pm0.27$  mV (n=10) for endocardial  $I_{Na}$  was obtained.

As demonstrated in Figure 4C-D, NS5806 induced a highly significant left-shift in midinactivation potential  $V_{\frac{1}{2}}$  for  $I_{Na}$ . For atrial  $I_{Na}$ , from  $V_{\frac{1}{2}} = -82.6\pm0.1$  mV to  $-85.1\pm0.1$  mV (n=8) and for Endo  $I_{Na}$  from  $V_{\frac{1}{2}} = -73.9\pm0.27$  mV to  $-77.3\pm0.21$  mV (n=9). As atrial cells have a more negative  $V_{\frac{1}{2}}$  compared to Endo cells, this suggest that as cells depolarize, atrial cells would lose excitability at less negative voltages than Endo cells. This tendency would be further enhanced by NS5806 suggesting that the effect of NS5806 on  $I_{Na}$  could dependent on resting membrane potential.

We next tested the effect of 10  $\mu$ M NS5806 on ventricular and atrial peak I<sub>Na</sub> using holding potentials ranging from -80 to -120 mV as illustrated by the representative atrial I<sub>Na</sub> recordings (Figure 5A). The current-voltage relationships for atrial (Figure 5B) and Endo I<sub>Na</sub> (Figure 5C) recorded at different holding potentials. For both cells types, I<sub>Na</sub> peak currents were smaller when the holding potential was -80 mV versus -120 mV. Application of NS5806 caused a progressive inhibition on I<sub>Na</sub> and the effect was more pronounced in atrial cells.

The differential effects of NS5806 on atrial and ventricular  $I_{to}$  and  $I_{Na}$  suggest different macromolecular ion channel complexes. To determine the expression level of mRNA encoding proteins thought to mediate I<sub>Na</sub> or I<sub>to</sub> as well as other K<sup>+</sup> currents, we performed qPCR experiments on mRNA isolated from Epi, Mid and Endo as well as from PM and right atrial appendage (App) (Figure 6). We observed a KChIP2 gradient across the right ventricle but no KCND3 gradient (encoding  $K_v$ 4.3). These observations suggest an I<sub>to</sub> gradient is present in the right ventricle, similar to what has been observed in the left ventricle. The relative mRNA expression levels of KCND3, showed an almost 4-fold higher expression in RV compared to RA. In contrast KCNA4 and KCNA5 encoding Kv1.4 and Kv1.5, respectively had a 2 fold higher expression in the atria. KCNH2 encoding Kv11.1 and KCNQ1 encoding K<sub>v</sub>7.1 was expressed at similar levels. We also tested the expression of several ancillary subunits and found a high expression of KCNIP2 encoding KChIP2, and a moderate expression of KCNE1, KCNE3 and KCNE4. For KCNIP2, the expression level in the atria was more than 2-fold higher than in ventricle and similar was found for KCNE4. A transmural expression gradient in the RV was also found for KCNE1. We observed no differences in any mRNA levels between PM and App. However, previous studies have shown that the electrophysiology is not uniform throughout the atria<sup>8,24</sup> suggesting some differences in ion channel expression throughout the atria. The expression level of the genes encoding proteins suspected to mediate INa was not significantly different between atria and ventricle, nor was any transmural gradient in expression found (Figure 6B)

## DISCUSSION

In the present work, we compared the effect of NS5806 on  $I_{to}$  and  $I_{Na}$  in canine atria and ventricle. In canine right ventricular wedges, phase 1 repolarization was increased in Epi while repolarization of Endo was largely unaffected. These changes resulted in an augmented J-wave on the ECG. In contrast, in atrial preparations, phase 1 repolarization was unaffected by NS5806 but excitability was reduced suggesting Na<sup>+</sup> channel block.

## Effect of NS5806 on Ito

It is believed that  $K_v4.3$  channels comprise the majority of transient outward K<sup>+</sup> channels in canine ventricle<sup>11,12</sup>. Several  $\beta$ -subunits can modify  $IK_v4.3$ , such as KChIP2 that increases peak  $IK_v4.3$  density, slows decay of the current and accelerates recovery from inactivation<sup>13</sup>. In the left and right ventricle, previous qPCR studies have shown that the  $I_{to}$  gradient across the ventricles is due to a KChIP2 gradient<sup>12</sup>. This gradient in mRNA was also confirmed at the protein level<sup>15</sup>. In the present study, qPCR analysis revealed a 3-fold lower expression of  $K_v4.3$  mRNA in atria compared to ventricle; interestingly, the expression of KChIP2 mRNA was 2-fold greater than RV Epi levels. Voltage clamp analysis of  $I_{to}$  showed that  $I_{to}$  was smaller in atria, in agreement with the qPCR data. However, this study raises an important question, namely what is the functional role of excess KChIP2 in canine atria? KChIP2 was initially identified as a  $\beta$ -subunit for  $K_v4$  channels; however, KChIP2 has been found to associate with  $I_{CaL}^{25}$  and  $I_{Na}^{26}$  suggesting it may alter the biophysical properties of many ion channels.

Epi I<sub>to</sub> was larger than atrial I<sub>to</sub> (Figure 2), however atrial cells had a prominent sustained current, I<sub>Kur</sub> that was not observed in Epi cells. NS5806 induced an increase in I<sub>to</sub> in both cells types, but the increase was more pronounced for Epi than atrial I<sub>to</sub>. The sustained current was unaffected by NS5806. Using qPCR, we found higher expression of K<sub>v</sub>1.4 and K<sub>v</sub>1.5 mRNA in the atria compared to ventricle. K<sub>v</sub>1.4 is thought to mediate a slowly recovering I<sub>to</sub>, I<sub>to</sub>, slow and K<sub>v</sub>1.5 is thought to mediate I<sub>Kur</sub>. For heterologously expressed K<sub>v</sub>1.5, 10  $\mu$ M NS5806 induced a minor block<sup>27</sup> consistent with a lack of effect on canine atrial sustained current. For heterologously expressed K<sub>v</sub>1.4, application of 10  $\mu$ M NS5806 resulted in an almost complete block <sup>27</sup>. The higher expression of K<sub>v</sub>1.4 and K<sub>v</sub>1.5 in the atria compared to the ventricle could explain why the relative increase in peak current induced by NS5806 was smaller in the atria than in Epi as NS5806 would enhance a K<sub>v</sub>4.3 mediated current component, block a K<sub>v</sub>1.4 component of I<sub>to</sub>, and leave a Kv1.5 mediated sustained current component unaffected.

The effect on different kinetic parameters of atrial  $I_{to}$  was addressed. Compared to ventricular  $I_{to}$ , atrial  $I_{to}$  had a slower current decay (Figure 2C,F and I). Application of NS5806 further slowed the current decay, similarly to the effects on ventricular I to. In a previous study, we found that NS5806 only slowed current decay of  $K_v4.3$  channels when co-expressed with KChIP2<sup>22,27</sup>, suggesting that at least part of the atrial  $I_{to}$  is mediated by  $K_v4.3$  in complex with KChIP2. The steady-state mid-inactivation potential for atrial  $I_{to}$  was shifted to more negative potentials by NS5806 similar to results seen in ventricular cells<sup>15</sup>.

## Effect of NS5806 on INa

Canine  $I_{Na}$  is functionally distinct in different areas of the heart. In the ventricles, Epi  $I_{Na}$  has a significant more negative steady-state mid-inactivation potential compared to Endo<sup>28</sup>. In agreement with other studies<sup>8,10,29,30</sup> we found that atrial  $I_{Na}$  has a higher current density and a more negative steady-state mid-inactivation potential compared to ventricular Endo  $I_{Na}$ . The negative steady-state mid-inactivation potential will tend to leave a larger fraction of atrial  $I_{Na}$  channels in an inactivated state. NS5806 produced a significant shift of steady-

J Cardiovasc Electrophysiol. Author manuscript; available in PMC 2012 September 1.

state inactivation to more negative potentials for both atrial and ventricular cells; this would be expected to lead to a larger pool of inactivated channels. The more negative midinactivation potential and more positive resting membrane potential in the atria (Table 1), can account for the greater depression of  $I_{Na}$  in atria vs. ventricles. We investigated the level of expression of  $Na_v 1.5$  subunits in the canine atria and in the three ventricular layers but did not detect significant differences in expression level of mRNA encoding  $Na_v 1.5$  or  $Na_v \beta 1-4$ suggesting mRNA levels measured by qPCR may not reflect protein levels in the membrane, or other proteins and regulatory factors account for the distinct behaviour of the channels.

#### Effect on action potential waveform

In RV ventricular wedges NS5806 (10  $\mu$ M) increased phase 1 repolarization in the Epi with no effect on repolarization of Endo as previously described<sup>22</sup>. In contrast there was no effect on phase 1 repolarization in atrial APs. Surprisingly, NS5806 reduced the AP amplitude, dV/dt and significantly depressed excitability in atrial tissue. NS5806 exerted similar effects in isolated atrial myocytes, whereas the effect on upstroke velocity of APs recoded from isolated ventricular Epi and Endo cells was minor. This disparate effect in atria vs. ventricle is likely due to the greater effect of NS5806 to depress I<sub>Na</sub> for reasons discussed above.

#### Physiological implications of Ito activation and INa block by NS5806

Many antiarrhythmic agents are effective in terminating atrial fibrillation in both the experimental and clinical setting due to their  $I_{Na}$  blocking ability. A major limitation of some of these agents is their potential proarrhythmic actions in the ventricle. Previously, it was shown that the antianginal agent, ranolazine was effective at suppressing atrial fibrillation with little effect in the ventricle<sup>8</sup>. This atrial effectiveness of ranolazine was due to intrinsic differences in sodium channels between atria and ventricle, namely the more negative steady-state inactivation curve<sup>8</sup>. In the present study, we also found intrinsic differences in sodium channels between atria and ventricle which resulted in greater sodium current inhibition by NS5806 in atria than in ventricle. While the effectiveness of NS5806 in terminating atrial fibrillation was not addressed in the present study, it is clear that NS5806 also exhibits atrial selectiveness in Na<sup>+</sup> channel blockade.

Of interest is the mechanism by which NS5806 activates  $I_{to}$ . We have previously shown this compound works through KChIP to cause an increase in the size of  $I_{to}^{15}$ . In the canine ventricle where  $K_v4.3$  levels are equivalent in Epi, Mid and Endo layers, the  $I_{to}$  enhancement was greatest in Epi and Mid cells where KChIP levels are highest. Although atrial tissue has high KChIP levels,  $K_v4.3$  levels were 3 times lower which may explain the modest increase in  $I_{to}$  following application of NS5806 in this tissue type, together with a potential block of atrial IK v1.4 and IKv1.5. In the uncompromised myocardium, application of this compound to ventricular wedges resulted in loss of the epicardial action potential dome and a Brugada-like ECG<sup>22</sup>. However, in pathophysiological condition such as heart failure, a reduction in  $I_{to}$  due to loss of both  $K_v4.3$  and KChIP levels has been observed<sup>31,32</sup>. If  $K_v4.3$  and KChIP levels are not severely diminished in heart failure, application of NS5806 may restore  $I_{to}$  and restore the spike and dome morphology.

The influence of action potential waveform and duration on excitation-contraction coupling is well known (for review see Sah et al.,  $2003^{33}$ ). The presence of a spike and dome morphology results in a significant increase in peak  $I_{CaL}$  as well as total charge when compared to a cell type where  $I_{to}$  and spike and dome morphology is absent<sup>34,35</sup>. These observations suggest the spike and dome morphology of the canine Epi action potential is a more efficient releaser of  $Ca^{2+}$  from the SR. In the setting of heart failure, where  $I_{to}$  is known to be reduced, this reduction in  $I_{to}$  would result in reduced  $Ca^{2+}$  influx into the cell via  $I_{CaL}$ . Augmentation of  $I_{to}$  in the setting of heart failure may have a beneficial effect on

cardiac output by restoring the spike-and-dome morphology resulting in increased  $Ca^{2+}$  influx during the course of an action potential.

In summary, NS5806 produces a prominent augmentation of  $I_{to}$  with little effect on  $I_{Na}$  in canine ventricles, but a potent inhibition of  $I_{Na}$  with little augmentation of  $I_{to}$  in canine atria. Molecular analysis of mRNA for atria and ventricle revealed more KChIP2,  $K_v 1.5$  and  $K_v 1.4$  and less  $K_v 4.3$  in atria versus ventricle

## Acknowledgments

**Financial Support:** This work was supported by Eva and Henry Fraenkel foundation [KC], the American Health Assistance Foundation [JMC]; the Danish National Research Foundation [SPO]; the National Institutes of Health [HL47678 to CA] and the Masons of New York State and Florida.

We are grateful to Judy Hefferon, Art Iodice, Robert Goodrow, and Trine Christensen for excellent technical assistance and NeuroSearch A/S for providing NS5806.

## Abbreviations

I <sub>to</sub>	transient outward current		
AP	action potential		
Ері	epicardium		
Endo	endocardium		
KChIP2	K Channel Interacting Protein 2		
DPP6	Dipeptidyl aminopeptidase-like protein 6		
PM	pectinate muscle		
СТ	crista terminalis		
RV	right ventricle		
LV	left ventricle		
RA	Right Atrial		

## References

- 1. Hume JR, Uehara A. Ionic basis of the different action potential configurations of single guinea-pig atrial and ventricular myocytes. J Physiol (Lond ). 1985; 368:525–44. [PubMed: 2416918]
- Giles WR, Imaizumi Y. Comparison of potassium currents in rabbit atrial and ventricular cells. J Physiol (Lond ). 1988; 405:123–45. [PubMed: 2855639]
- Sridhar A, da Cunha DN, Lacombe VA, Zhou Q, Fox JJ, Hamlin RL, Carnes CA. The plateau outward current in canine ventricle, sensitive to 4-aminopyridine, is a constitutive contributor to ventricular repolarization. Br J Pharmacol. 2007; 152(6):870–9. [PubMed: 17700726]
- Fedida D, Wible BA, Wang ZG, Fermini B, Faust F, Nattel S, Brown AM. Identity of a novel delayed rectifier current from human heart with a cloned K <sup>+</sup> channel current. Circ Res. 1993; 73:210–6. [PubMed: 8508531]
- Melnyk P, Zhang L, Shrier A, Nattel S. Differential distribution of Kir2.1 and Kir2.3 subunits in canine atrium and ventricle. Am J Physiol Heart Circ Physiol. 2002; 283(3):H1123–H1133. [PubMed: 12181143]
- Fedida D, Eldstrom J, Hesketh JC, Lamorgese M, Castel L, Steele DF, Van Wagoner DR. Kv1.5 is an important component of repolarizing K<sup>+</sup> current in canine atrial myocytes. Circ Res. 2003; 93(8): 744–51. [PubMed: 14500335]

- Dhamoon AS, Pandit SV, Sarmast F, Parisian KR, Guha P, Li Y, Bagwe S, Taffet SM, Anumonwo JM. Unique Kir2.x properties determine regional and species differences in the cardiac inward rectifier K<sup>+</sup> current. Circ Res. 2004; 94(10):1332–9. [PubMed: 15087421]
- Burashnikov A, Di Diego JM, Zygmunt AC, Belardinelli L, Antzelevitch C. Atrium-selective sodium channel block as a strategy for suppression of atrial fibrillation: differences in sodium channel inactivation between atria and ventricles and the role of ranolazine. Circulation. 2007; 116(13):1449–57. [PubMed: 17785620]
- Furukawa T, Koumi S, Sakakibara Y, Singer DH, Jia H, Arentzen CE, Backer CL, Wasserstrom JA. An analysis of lidocaine block of sodium current in isolated human atrial and ventricular myocytes. J Mol Cell Cardiol. 1995; 27(2):831–46. [PubMed: 7776390]
- Li GR, Lau CP, Shrier A. Heterogeneity of sodium current in atrial vs epicardial ventricular myocytes of adult guinea pig hearts. J Mol Cell Cardiol. 2002; 34(9):1185–94. [PubMed: 12392892]
- Dixon EJ, Shi W, Wang H-S, McDonald C, Yu H, Wymore RS, Cohen IS, McKinnon D. Role of the Kv4.3 K+ channel in ventricular muscle. A molecular correlate for the transient outward current. Circ Res. 1996; 79:659–68. [PubMed: 8831489]
- Rosati B, Grau F, Rodriguez S, Li H, Nerbonne JM, McKinnon D. Concordant expression of KChIP2 mRNA, protein and transient outward current throughout the canine ventricle. J Physiol. 2003; 548(Pt 3):815–22. [PubMed: 12598586]
- Deschênes I, DiSilvestre D, Juang GJ, Wu RC, An WF, Tomaselli GF. Regulation of K<sub>v</sub>4.3 current by KChIP2 splice variants: a component of native cardiac I<sub>to</sub>? Circulation. 2002; 106(4):423–9. [PubMed: 12135940]
- 14. Rosati B, Pan Z, Lypen S, Wang HS, Cohen I, Dixon JE, McKinnon D. Regulation of KChIP2 potassium channel beta subunit gene expression underlies the gradient of transient outward current in canine and human ventricle. J Physiol. 2001; 533(Pt 1):119–25. [PubMed: 11351020]
- Calloe K, Soltysinska E, Jespersen T, Lundby A, Antzelevitch C, Olesen SP, Cordeiro JM. Differential effects of the transient outward K<sup>+</sup> current activator NS5806 in the canine left ventricle. J Mol Cell Cardiol. 2010; 48(1):191–200. [PubMed: 19632239]
- 16. Radicke S, Cotella D, Graf EM, Ravens U, Wettwer E. Expression and function of dipeptidylaminopeptidase-like protein 6 as a putative b-subunit of human cardiac transient outward current encoded by Kv4.3. J Physiol. 2005; 565(Pt 3):751–6. [PubMed: 15890703]
- Delpon E, Cordeiro JM, Nunez L, Thomsen PE, Guerchicoff A, Pollevick GD, Wu Y, Kanters JK, Larsen CT, Hofman-Bang J, Burashnikov E, Christiansen M, Antzelevitch C. Functional effects of KCNE3 mutation and its role in the development of Brugada syndrome. Circ Arrhythm Electrophysiol. 2008; 1(3):209–18. [PubMed: 19122847]
- Lundby A, Olesen SP. KCNE3 is an inhibitory subunit of the Kv4.3 potassium channel. BBRC. 2006; 346(3):958–67. [PubMed: 16782062]
- Akar FG, Wu RC, Deschênes I, Armoundas AA, Piacentino V III, Houser SR, Tomaselli GF. Phenotypic differences in transient outward K<sup>+</sup> current of human and canine ventricular myocytes: insights into molecular composition of ventricular I<sub>to</sub>. Am J Physiol Heart Circ Physiol. 2004; 286(2):H602–H609. [PubMed: 14527940]
- 20. Calloe K, Chlus N, Nof E, Jespersen T, Olesen SP, Antzelevitch C, Cordeiro JM. Comparison of the effects of the transient outward potassium channel activator NS5806 on canine atrial and ventricular cardiomyocytes. Biophys J. 2010; 98:334a. Ref Type: Abstract.
- Dumaine R, Cordeiro JM. Comparison of K+ currents in cardiac Purkinje cells isolated from rabbit and dog. J Mol Cell Cardiol. 2007; 42(2):378–89. [PubMed: 17184792]
- 22. Calloe K, Cordeiro JM, Di Diego JM, Hansen RS, Grunnet M, Olesen SP, Antzelevitch C. A transient outward potassium current activator recapitulates the electrocardiographic manifestations of Brugada syndrome. Cardiovasc Res. 2009; 81(4):686–94. [PubMed: 19073629]
- Yue L, Feng J, Li GR, Nattel S. Characterization of an ultrarapid delayed rectifier potassium channel involved in canine atrial repolarization. J Physiol. 1996; 496 (Pt 3):647–62. [PubMed: 8930833]
- 24. Feng J, Yue L, Wang Z, Nattel S. Ionic mechanisms of regional action potential heterogeneity in the canine right atrium. Circ Res. 1998; 83(5):541–51. [PubMed: 9734477]

- Page 11
- 25. Thomsen MB, Wang C, Ozgen N, Wang HG, Rosen MR, Pitt GS. Accessory subunit KChIP2 modulates the cardiac L-type calcium current. Circ Res. 2009; 104(12):1382–9. [PubMed: 19461043]
- 26. Deschenes I, Armoundas AA, Jones SP, Tomaselli GF. Post-transcriptional gene silencing of KChIP2 and Navbeta1 in neonatal rat cardiac myocytes reveals a functional association between Na and Ito currents. J Mol Cell Cardiol. 2008; 45(3):336–46. [PubMed: 18565539]
- Lundby A, Jespersen T, Schmitt N, Grunnet M, Olesen SP, Cordeiro JM, Calloe K. Effect of the I(to) activator NS5806 on cloned K(V)4 channels depends on the accessory protein KChIP2. Br J Pharmacol. 2010; 160(8):2028–44. [PubMed: 20649599]
- Cordeiro JM, Mazza M, Goodrow R, Ulahannan N, Antzelevitch C, Di Diego JM. Functionally distinct sodium channels in ventricular epicardial and endocardial cells contribute to a greater sensitivity of the epicardium to electrical depression. Am J Physiol Heart Circ Physiol. 2008; 295(1):H154–H162. [PubMed: 18456729]
- Sakakibara Y, Wasserstrom JA, Furukawa T, Jia H, Arentzen CE, Hartz RS, Singer DH. Characterization of the sodium current in single human atrial myocytes. Circ Res. 1992; 71:535– 46. [PubMed: 1323431]
- Sakakibara Y, Furukawa T, Singer DH, Jia H, Backer CL, Arentzen CE, Wasserstrom JA. Sodium current in isolated human ventricular myocytes. Am J Physiol. 1993; 265(4 Pt 2):H1301–H1309. [PubMed: 8238418]
- 31. Akar FG, Wu RC, Juang GJ, Tian Y, Burysek M, DiSilvestre D, Xiong W, Armoundas AA, Tomaselli GF. Molecular mechanisms underlying K+ current downregulation in canine tachycardia-induced heart failure. Am J Physiol Heart Circ Physiol. 2005; 288(6):H2887–H2896. [PubMed: 15681701]
- 32. Zicha S, Xiao L, Stafford S, Cha TJ, Han W, Varro A, Nattel S. Transmural expression of transient outward potassium current subunits in normal and failing canine and human hearts. J Physiol. 2004; 561(Pt 3):735–48. [PubMed: 15498806]
- 33. Sah R, Ramirez RJ, Oudit GY, Gidrewicz D, Trivieri MG, Zobel C, Backx PH. Regulation of cardiac excitation-contraction coupling by action potential repolarization: role of the transient outward potassium current (I(to)). J Physiol. 2003; 546(Pt 1):5–18. [PubMed: 12509475]
- 34. Cordeiro JM, Greene L, Heilmann C, Antzelevitch D, Antzelevitch C. Transmural heterogeneity of calcium activity and mechanical function in the canine left ventricle. Am J Physiol Heart Circ Physiol. 2004; 286(4):H1471–H1479. [PubMed: 14670817]
- Hund TJ, Rudy Y. Rate dependence and regulation of action potential and calcium transient in a canine cardiac ventricular cell model. Circulation. 2004; 110(20):3168–74. [PubMed: 15505083]

Calloe et al.



#### Figure 1.

Effect of NS5806 on arterially perfused atrial and ventricular preparations paced at BCL=1 sec. **Panel A**) Representative action potentials (AP) recordings from right atrial pectinate muscle (PM) preparation in presence of 0–15  $\mu$ M NS5806 (n=4). **Panel B**) Representative AP recordings and corresponding ECG from a right ventricular wedge in absence and presence of 10  $\mu$ M NS5806 (n=5). **Panel C**) Representative APs and corresponding dV/dt recorded from isolated canine cardiomyocytes at a BCL of 1 sec. APs recorded from a atrial cell (left panel), LV Epi cell (middle panel), and LV Endo cell (left panel) in absence (black traces) and presence of 10  $\mu$ M NS5806 (gray traces). The APs recorded from the LV Epi cell were 8 and 24 sec after application of 10  $\mu$ M NS5806. The results are summarized in Table 1.

Calloe et al.



#### Figure 2.

 $I_{to}$  in canine atrial and ventricular cells in absence and presence of NS5806 (10  $\mu$ M). Currents were activated by the voltage clamp protocol (top of figure) and Cd<sup>2+</sup> was present to block I<sub>CaL</sub>. **Panel A**) Representative I<sub>to</sub> traces recorded from canine atrial cells. **Panel B**) Mean I–V relation for atrial peak I<sub>to</sub> as well as for the sustained component **Panel C**) I<sub>to</sub> decay as a function of voltage under control (n=14). **Panel D**) Representative I<sub>to</sub> traces recorded from canine LV Endo cells. **Panel E**) Mean I–V relation for peak I<sub>to</sub>. **Panel F**) I<sub>to</sub> decay as a function of voltage (n=9). **Panel G**) Representative I<sub>to</sub> traces recorded from canine LV Epi cells. **Panel H**) Mean I–V relation for peak I<sub>to</sub>. **Panel I**) I<sub>to</sub> decay as a function of voltage (n=10).



#### Figure 3.

Voltage dependence of inactivation of atrial  $I_{to}$ . Representative traces recorded under control conditions (**Panel A**) and after application of 10  $\mu$ M NS5806 (**Panel B**) showing voltage dependence of inactivation of  $I_{to}$ . **Panel C**) The currents were normalized to max-current and plotted as a function of the conditioning pulse and Boltzmann curves were fit to the data. In the absence of drug, the mid-inactivation voltage was  $V_{1/2}$ =-33.8±0.46 mV and the slope factor k=-7.15±0.40. In presence of NS5806 V<sub>1/2</sub> was -41.1±0.50 mV and k= -7.35±0.42 (n=8). **Panel D**) Time-dependence of recovery from inactivation of atrial I<sub>to</sub>. Currents were elicited by the depicted protocol and normalized recovered currents were plotted as a function of the interpulse interval (n=6).



## Figure 4.

I<sub>Na</sub> recorded from canine atrial cells or endocardial cells. Cd<sup>2+</sup> was present to block I<sub>CaL</sub>. **Panel A)** Representative I<sub>Na</sub> traces recorded from atrial (n=8) and Endo cells (n=9) and corresponding mean I–V relation for peak current is shown. **Panel B)** Representative steady-state inactivation recordings from an atrial or Endo cell. The peak-current following a 500 ms prepulse was normalized to the maximum current and plotted against the conditioning potential to obtain the availability of channels. A Boltzmann equation was fit to the data resulting in a mid-inactivation (V<sub>1/2</sub>) of  $-82.6\pm0.12$  mV (n=8) for atrial and  $-73.9\pm0.27$  mV (n=9) for Endo I <sub>Na</sub>. **Panel C**) Steady-state inactivation for I<sub>Na</sub> recorded from atrial cells in absence and presence of 10 µM NS5806. V<sub>1/2</sub> =  $-82.6\pm0.1$  mV in control and  $-85.1\pm0.1$  mV in control and  $-73.9\pm0.27$  mV in presence of NS5806 (n=8). **Panel D**) Steady-state inactivation for I<sub>Na</sub> recorded from isolated LV Endo cells in absence and presence of 10 µM NS5806. N<sub>1/2</sub> =  $-73.9\pm0.27$  mV in control and  $-77.3\pm0.21$  mV in presence of NS5806 (n=9).

Calloe et al.



#### Figure 5.

The effect of different holding potentials on atrial and ventricular I  $_{Na}$  in absence and presence of 10  $\mu$ M NS5806. The voltage clamp protocol is shown at the top of the figure and Cd<sup>2+</sup> was present to block I<sub>CaL</sub>. **Panel A**) Representative I<sub>Na</sub> traces recorded from canine right atrial cells using holding potentials of either -80 mV, -100 mV or -120 mV. **Panel B**) Mean I–V relation for atrial I<sub>Na</sub> peak current recorded using different holding potentials in absence and presence of 10  $\mu$ M NS5806 (n=5). **Panel C**) Mean I–V relation for LV Endo I<sub>Na</sub> peak current recorded using different holding potentials in absence and presence of 10  $\mu$ M NS5806 (n=5).

Calloe et al.



#### Figure 6.

qPCR of canine right ventricular (Epi, Mid, and Endo) and right atrial pectinate muscle (PM) and appendage (App) tissue isolated from 5 canine hearts. Taqman based assays (Applied Biosystems) were used to quantify mRNA of interest which was normalized to *cyclophilin B* expression. **Panel A**) Expression of K<sup>+</sup> channel subunit mRNA. **Panel B**) Expression of Na<sup>+</sup> channel subunit mRNA.

## Table 1

Action potentials were recorded from isolated canine atrial or left ventricular Epi or Endo cells at a basic cycle length (BCL) of 1 sec in the absence or presence of 10  $\mu$ M NS5806. The effects of NS5806 on upstroke velocity (dV/dt), on phase 0 (Ph0), phase 1(Ph1) and phase 2 (Ph2) as well as on action potential duration (ADP) and on resting membrane potential (E<sub>rest</sub>) of different cell types are summarized.

	Atria	Ері	Endo
ΔdV/dt	-71±7%, (n=6)***	-9.8±4.8%, (n=5) *	-4.9±2.8%, (n=4) *
Ph0 Amplitude + NS5806	108±4 mV	124±3 mV	126±8 mV
	72±5 mV, (n=6) **	124±3 mV, (n=5), NS	125±7, (n=4), NS
Notch, Ph1% of Ph2 + NS5806	-	28.2±11.1%, (n=5) no Ph2 in 5/5	5.5±1.0% 5.1±1.8%, (n=4), NS
APD90 + NS5806	294±28 ms	527±46 mS	535±56 mS
	187±24 ms, (n=6) ***	59±6 mS, (n=5) ***	450±28 mS, (n=4), NS
E <sub>rest</sub> + NS5806	-71.9±0.3 mV	-87.1±5 mV	-86.0±6 mV
	-72.6±0.6 mV, (n=6), NS	-87.4±3 mV, (n=5), NS	-84.9±6 mV, (n=4), NS