

# Adenosine Triphosphate-Sensitive Potassium Channel Kir Subunits Implicated in Cardioprotection by Diazoxide

Matthew C. Henn, MD; M. Burhan Janjua, MD; Evelyn M. Kanter, BS; Carol M. Makepeace, BS; Richard B. Schuessler, PhD; Colin G. Nichols, PhD; Jennifer S. Lawton, MD

**Background**—ATP-sensitive potassium ( $K_{ATP}$ ) channel openers provide cardioprotection in multiple models. Ion flux at an unidentified mitochondrial  $K_{ATP}$  channel has been proposed as the mechanism. The renal outer medullary kidney potassium channel subunit, potassium inward rectifying (Kir)1.1, has been implicated as a mitochondrial channel pore-forming subunit. We hypothesized that subunit Kir1.1 is involved in cardioprotection (maintenance of volume homeostasis and contractility) of the  $K_{ATP}$  channel opener diazoxide (DZX) during stress (exposure to hyperkalemic cardioplegia [CPG]) at the myocyte and mitochondrial levels.

**Methods and Results**—Kir subunit inhibitor Tertiapin Q (TPN-Q) was utilized to evaluate response to stress. Mouse ventricular mitochondrial volume was measured in the following groups: isolation buffer; 200  $\mu$ mol/L of ATP; 100  $\mu$ mol/L of DZX+200  $\mu$ mol/L of ATP; or 100  $\mu$ mol/L of DZX+200  $\mu$ mol/L of ATP+TPN-Q (500 or 100 nmol/L). Myocytes were exposed to Tyrode's solution (5 minutes), test solution (Tyrode's, cardioplegia [CPG], CPG+DZX, CPG+DZX+TPN-Q, Tyrode's+TPN-Q, or CPG+TPN-Q), N=12 for all (10 minutes); followed by Tyrode's (5 minutes). Volumes were compared. TPN-Q, with or without DZX, did not alter mitochondrial or myocyte volume. Stress (CPG) resulted in myocyte swelling and reduced contractility that was prevented by DZX. TPN-Q prevented the cardioprotection afforded by DZX (volume homeostasis and maintenance of contractility).

**Conclusions**—TPN-Q inhibited myocyte cardioprotection provided by DZX during stress; however, it did not alter mitochondrial volume. Because TPN-Q inhibits Kir1.1, Kir3.1, and Kir3.4, these data support that any of these Kir subunits could be involved in the cardioprotection afforded by diazoxide. However, these data suggest that mitochondrial swelling by diazoxide does not involve Kir1.1, 3.1, or 3.4. (*J Am Heart Assoc.* 2015;4:e002016 doi: 10.1161/JAHA.115.002016)

**Key Words:** cardioplegia • ion channels • myocardial stunning • potassium

In an isolated model of myocardial stunning, cardiac myocytes demonstrate significant swelling and reduced contractility during exposure to 3 different stresses: hypothermic hyperkalemic cardioplegia (CPG), hypo-osmotic stress, or metabolic inhibition.<sup>1–3</sup> The ATP-sensitive potassium channel ( $K_{ATP}$ ) opener, diazoxide (DZX), ameliorates these stress-induced derangements in mouse, rabbit, and

human tissue.<sup>1–5</sup> A role of  $K_{ATP}$  channels in ischemic preconditioning, by coupling membrane conductance of  $K^+$  ions to the metabolic state of the cell, has also been well defined in multiple animal models.<sup>6–10</sup> However, no precise mechanism of cardioprotection has been identified.

$K_{ATP}$  channels are formed as hetero-octamers of subunits that include 4 potassium inward rectifying (Kir) subunits and 4 regulatory subunits from the sulfonylurea receptor family (SUR1/2).<sup>11–13</sup> Different combinations of Kir and SUR isoforms exist in different tissues, each with unique properties and function. For example, the predominant combination of subunits found in murine ventricle is Kir6.2 and SUR2A, whereas murine atria mainly consist of Kir6.2 and SUR1.<sup>14</sup> Identification of these subunits has allowed for genetic manipulation to more specifically evaluate the mechanism of DZX.

Using  $K_{ATP}$  subunit-knockout animal models, we have shown that the cardioprotection afforded by DZX requires the regulatory subunit, SUR1, but the pore-forming subunit involved remains elusive.<sup>15,16</sup> We have also demonstrated that the sarcolemmal  $K_{ATP}$  channel (s $K_{ATP}$ ) is implicated in

From the Division of Cardiothoracic Surgery, Departments of Surgery (M.C.H., M.B.J., E.M.K., C.M.M., R.B.S., J.S.L.) and Cell Biology and Physiology (C.G.N.), Washington University School of Medicine, St. Louis, MO.

Presented at the American Heart Association Scientific Sessions, November 15–19, 2014 in Chicago, IL.

**Correspondence to:** Jennifer S. Lawton, MD, Washington University School of Medicine, 660 S. Euclid Ave, Campus Box 8234, St. Louis, MO 63110.

E-mail: lawtonj@wustl.edu

Received March 24, 2015; accepted April 20, 2015.

© 2015 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley Blackwell. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

myocyte swelling secondary to stress, and that knockout of the pore-forming subunit of this channel (Kir6.2) provides resistance to myocyte swelling.<sup>2,17</sup> However, there is evidence that DZX cardioprotection does not require the  $sK_{ATP}$  channel, leading us to propose mechanisms that may include non- $K_{ATP}$  channel or mitochondrial  $K_{ATP}$  channel ( $mK_{ATP}$ ) locations of action.<sup>15,16,18–21</sup>

Regulation of mitochondrial volume through a purported  $mK_{ATP}$  channel is one of the proposed mechanisms of action of DZX.<sup>22–24</sup> Because no molecular composition of  $mK_{ATP}$  has been determined, identification of  $mK_{ATP}$  channel subunits has to date involved indirect methods, such as pharmacological inhibition or observed changes in mitochondrial volume (proposed assay of  $mK_{ATP}$  activity).<sup>24,25</sup> Using such strategies, the inward rectifier potassium channel subunit, Kir1.1 (renal outer medullary kidney channel; ROMK) has been implicated as a pore-forming subunit of the  $mK_{ATP}$  channel utilizing the nonspecific inhibitor Tertiapin Q (TPN-Q).<sup>24</sup>

The present study was conducted to assess potential roles of Kir1.1, Kir3.1, and Kir3.4 in the cardioprotection of DZX at the cellular and mitochondrial levels using the Kir subunit inhibitor, TPN-Q.

## Methods

All animal procedures were approved by the Animal Studies Committee at Washington University School of Medicine, and all animals received humane care in compliance with the National Institute of Health's (NIH) Guide to Care and Use of Laboratory Animals.<sup>26</sup>

## Mitochondrial Isolation

Mice (both sexes, 6 weeks to 5 months and 15 to 30 g in weight) were anesthetized with 3% Avertin (0.3 g of 2,2,2-tribromoethanol, 0.186 mL of 2-methyl-2-butanol, and 9.814 mL of sterile water) intraperitoneally, and rapid cardiectomy was performed. To isolate mitochondria, atria were excised and discarded whereas ventricular tissue was rapidly minced in cold buffer (in mmol/L: 10 HEPES (*N*-[2-hydroxyethyl]<sup>1,3–5</sup> piperazine-*N*-[4-butanedisulfonic acid]), 1 EDTA-potassium, 250 sucrose, adjusted to a pH of 7.1 with 20% KOH and transferred to a 10-mL glass tube with a Teflon pestle (Glas-Col Homogenizer, Terre Haute, IN), and volume was adjusted to 7 mL with buffer. Tissue was mechanically homogenized with a Teflon pestle driven by a low-speed motor drive shaft set at 120 rpm. The homogenate was transferred to 6 microcentrifuge tubes and centrifuged at 900g for 10 minutes at 4°C. The supernatant was combined into a clean test tube and mixed to get a homogeneous solution that was divided equally between 6 clean microcentrifuge tubes

and centrifuged at 5000g for 15 minutes. One pellet was resuspended in 100  $\mu$ L of buffer, and 10  $\mu$ L was taken in duplicate for the Bradford protein assay (Thermo Scientific; Rockford, IL) to determine total protein. Each pellet was maintained on ice and was resuspended in test media volume to equal 0.3  $\mu$ g/ $\mu$ L in order to standardize protein content.

## Mitochondrial Volume Measurement

The volume of isolated mitochondria was measured after suspension in test solution: (1) isolation buffer (No ATP;  $n=13$ ); (2) 200  $\mu$ mol/L of ATP ( $n=12$ ); (3) 200  $\mu$ mol/L of ATP and 100  $\mu$ mol/L of DZX (Sigma-Aldrich, St. Louis, MO) ( $n=12$ ); (4) 200  $\mu$ mol/L of ATP, 100  $\mu$ mol/L of DZX, and 500 nmol/L of TPN-Q ( $n=7$ ); or (5) 200  $\mu$ mol/L of ATP, 100  $\mu$ mol/L of DZX, and 100 nmol/L of TPN-Q ( $n=6$ ). Isolation buffer (10 mmol/L of HEPES, 200 mmol/L of mannitol, 50 mmol/L of sucrose, 1 mmol/L of EGTA; pH 7.2) was used as a control solution. ATP has been shown to close mitochondrial  $K_{ATP}$  channels, so 200  $\mu$ mol/L of ATP was used to slow the initial rate of mitochondrial swelling (0% Mito  $K_{ATP}$  activity).<sup>22</sup> Conversely, DZX activates “Mito  $K_{ATP}$ ” channels, and 100  $\mu$ mol/L of DZX was added to achieve maximal activation of mitochondrial  $K_{ATP}$  channels (100% Mito  $K_{ATP}$  activity). We were unable to reproduce the TPN-Q dose-response relationships on mitochondrial volume demonstrated by other investigators using 4 TPN-Q concentrations (0.5, 10, 90, or 1000 pmol/L) despite attempts utilizing 3 different vehicles for TPN-Q: 20% acetonitrile, water, and HEPES.<sup>24</sup> In the present study, water was used as a vehicle for TPN-Q because of its stability in the medium and because 20% acetonitrile alone resulted in myocyte swelling owing to its cyanide moiety.

Mitochondrial matrix volume measurements were obtained using a light-scattering technique,<sup>27</sup> where the absorbance, at 520 nm, of a solution of isolated mitochondria was obtained every 14 seconds for the period of 3 minutes using UV Probe 2.33 (Shimadzu Scientific Instruments, Columbia, MD) and a spectrophotometer (UV-1700 Spectrophotometer; Shimadzu Scientific Instruments, Columbia, MD).

## Myocyte Isolation

Ventricular myocytes were isolated from C57BL/6J mice of either sex (age 6 weeks to 5 months and 15 to 30 g in weight), as previously described.<sup>15</sup> Mice were anesthetized with 2.5% Avertin intraperitoneally. Heparin (0.1 mL) was administered intraperitoneally. Rapid cardiectomy was performed and solution A (described below) was perfused through the aorta for 5 minutes. The heart was then perfused at 37°C for 12 to 20 minutes with solution B (described below). Ventricles were removed, minced, and placed into solution C (described below) and gently dispersed by glass pipette. Cells were allowed to

centrifuge by gravity, and serial washings were performed every 10 minutes for 15 to 20 minutes. Cells were used within 5 hours and randomized to treatment group. A typical yield of viable myocytes was 65% to 75%.

Solution A consisted of (in mmol/L, except as noted) 116 NaCl; 5.36 KCl; 0.97 Na<sub>2</sub>HPO<sub>4</sub>; 1.47 KH<sub>2</sub>PO<sub>4</sub>; 21.10 HEPES (*N*-[2-hydroxyethyl] piperazine-*N'*-[4-butanedisulfonic acid]); 11.65 glucose; 26.50  $\mu$ mol/L of phenol red (Sigma-Aldrich); 3.72 MgCl<sub>2</sub>; 4.40 NaHCO<sub>3</sub>; essential vitamins (100 $\times$ , 10 mL; GIBCO, Grand Island, NY); and amino acids (50 $\times$ , 20 mL; GIBCO). Solution B consisted of solution A plus 10  $\mu$ mol/L of CaCl<sub>2</sub>; 1.2 mg/mL of collagenase (Type 2; Worthington Biochemical Corporation, Freehold, NJ). Solution C consisted of solution A plus 5 mg/mL of BSA (Sigma-Aldrich); 1.25 mg/mL of taurine; and 150  $\mu$ mol/L of CaCl<sub>2</sub>.

Myocytes were exposed to 37°C control TYR for 5 minutes to obtain baseline volume. Any changes in cell volume secondary to the isolation would be evident during this period. Myocytes were then exposed to test solution (10 minutes) followed by re-exposure to TYR solution (5 minutes). Test solutions included the following groups (*n*=12 for each): (1) TYR; (2) CPG; (3) CPG+100  $\mu$ mol/L of DZX; (4) CPG+100  $\mu$ mol/L of DZX+TPN-Q 200 nm/L; (5) TYR+TPN-Q; and (6) CPG+TPN-Q.

CPG consisted of (in mmol/L): NaCl 110, NaHCO<sub>3</sub> 10, KCl 16, MgCl<sub>2</sub> 16, and CaCl<sub>2</sub> 1.2 and was equilibrated with 95% O<sub>2</sub> to 5% CO<sub>2</sub> and titrated to the pH of 7.3 with 10% NaHCO<sub>3</sub> solution. Diazoxide (7-chloro-3-methyl-1,2,4-benzothiadiazine-1,1-dioxide<sup>4</sup>; Sigma-Aldrich) dose of 100  $\mu$ mol/L was utilized because it was effective in ameliorating cell swelling secondary to stress in previous studies.<sup>1–4</sup> A stock solution of DZX was made by dissolving DZX in 0.1% dimethyl sulfoxide (DMSO), at which concentration DMSO has no effect on cell volume.<sup>28</sup>

## Myocyte Volume Measurement

Myocytes were used on the day of isolation and were not cultured. Myocytes were visualized on an inverted microscope stage (IX-51; Olympus, Tokyo, Japan), as previously described.<sup>15</sup> After 5 minutes, the chamber was perfused at a rate of 3 mL/min with TYR (in mmol/L): NaCl 130, KCl 5, CaCl<sub>2</sub> 2.5, MgSO<sub>4</sub> 1.2, NaHCO<sub>3</sub> 24, Na<sub>2</sub>HPO<sub>4</sub> 1.75, and glucose 10 (buffered to a pH of 7.4 using 95% O<sub>2</sub> to 5% CO<sub>2</sub>). After viability was confirmed, myocyte images were captured using video-based edge detection software (IonOptix, Milton, MA) and volume measured every 5 minutes, as previously described.<sup>5</sup>

## Myocyte Contractility

Myocyte contractility was measured using a video-based edge detection system (IonOptix). Cells were paced using a field stimulator (MyoPacer; IonOptix) at a voltage of 105 above

threshold at a frequency of 1 Hz with a 5-ms duration to avoid occurrence of fusion beats. After 5 minutes of stimulation, data were obtained from 12 to 30 consecutive beats and averaged. Parameters of contractility included percentage of cell shortening, maximal velocity of shortening, and percentage of cell relengthening, as previously described.<sup>5</sup> Contractility was measured at baseline and after 5 minutes of re-exposure to TYR. Cells that showed less than 7% cell shortening at baseline were excluded.

## Statistical Analysis

Data were analyzed using SYSTAT 13 (Systat Software, Inc, Point Richmond, CA). All data are presented as mean $\pm$ SEM relative to baseline. A repeated-measures ANOVA was used for sequential time-based measurements for each test solution against its own baseline and control values. Using Fisher's least significant different test, post-hoc multiple comparisons were done between different test groups at different time points during test solution and re-exposure periods. Probability values <0.05 were considered significant. A Shapiro-Wilk test was used to test for normality. If the data failed the normality test, a nonparametric (Friedman's nonparametric repeated-measures comparison) was used. Statistical analysis was performed using SyStat 10.2 (Systat Software).

## Results

### Mitochondrial Matrix Volume


Isolated mitochondrial volume is represented in Figure 1. There were no statistically significant differences in mitochondrial volume between any of the groups.

### Myocyte Volume

Myocyte volume over time is represented in Figure 2. Myocytes demonstrated no significant volume change from baseline during exposure to TYR with or without TPN-Q. Myocytes did demonstrate significant swelling when exposed to CPG (*P*<0.05 vs. TYR). This swelling was prevented by addition of DZX (*P*<0.05 vs. CPG). The further addition of TPN-Q prevented this benefit (volume homeostasis) of DZX (CPG+DZX+TPN-Q *P*<0.05 vs. CPG+DZX). As expected, the interaction between experimental groups and time was significant (*P*<0.001).


### Myocyte Contractility

Myocyte contractility is presented in Figure 3. Contractility remained unaltered in the TYR group. Myocytes demonstrated a significant decline in all 3 parameters of contractility after




**Figure 1.** Mitochondrial matrix volume. Isolated mouse mitochondria were exposed to: (1) isolation buffer (no ATP); (2) 200  $\mu\text{mol/L}$  of ATP; (3) 200  $\mu\text{mol/L}$  of ATP and 100  $\mu\text{mol/L}$  of DZX; (4) 200  $\mu\text{mol/L}$  of ATP, 100  $\mu\text{mol/L}$  of DZX, and 500 nmol/L of TPN-Q; or (5) 200  $\mu\text{mol/L}$  of ATP, 100  $\mu\text{mol/L}$  of DZX, and 100 nmol/L of TPN-Q and volume measured using light scattering (at 520 nm) for 2 minutes. Mitochondrial matrix volume is represented as 1/percent change in absorbance (mitochondrial matrix swelling is inversely related to absorbance measured at 520 nm) over time. Data are represented as mean $\pm$ SEM. DZX indicates diazoxide; TPN-Q, tertiapin Q.

exposure to CPG ( $P<0.05$  vs. TYR) that was worsened by the addition of TPN-Q. The addition of DZX to CPG prevented the significant reduction in contractility observed in the CPG group. TPN-Q inhibited the protection provided by DZX.



**Figure 2.** TPN-Q prevents beneficial maintenance of myocyte volume during stress by DZX. Isolated myocytes were exposed to control Tyrode's physiologic solution (TYR) for 5 minutes (time 0 to 5), stress (CPG, CPG+DZX, CPG+DZX+TPN-Q, CPG+TPN-Q, and TYR+TPN-Q) for 10 minutes (time 5 to 15), followed by TYR for 5 minutes (time 15 to 20). Myocyte volume was compared. Data are represented as mean $\pm$ SEM.  $*P<0.05$  vs. TYR;  $^{\Psi}P<0.05$  vs. CPG+DZX. CPG indicates hypothermic hyperkalemic cardioplegia; DZX, diazoxide; TPN-Q, tertiapin Q.



**Figure 3.** TPN-Q prevents beneficial maintenance of myocyte contractility after stress by DZX. Isolated mouse myocytes were exposed to control Tyrode's physiologic solution (TYR) for 5 minutes (time 0 to 5), stress (CPG, CPG+DZX, CPG+DZX+TPN-Q, CPG+TPN-Q, and TYR+TPN-Q) for 10 minutes (time 5 to 15), followed by TYR for 5 minutes (time 15 to 20). Myocyte contractility was compared. Data are represented as mean $\pm$ SEM.  $*P<0.05$  vs. TYR;  $^{\Psi}P<0.05$  vs. CPG+DZX. CPG indicates hypothermic hyperkalemic cardioplegia; DZX, diazoxide; TPN-Q, tertiapin Q.

## Discussion

In previous studies, we have shown that myocyte swelling and reduced contractility in response to stress (hyperkalemic CPG, hypoosmotic stress, and metabolic inhibition) are ameliorated by addition of the  $K_{ATP}$  channel opener, DZX.<sup>1–5</sup> Our efforts to localize the precise mechanism of cardioprotection afforded by DZX have implicated an involvement of the  $K_{ATP}$  channel, SUR1,<sup>16,19</sup> but not the Kir6.2 subunit.<sup>2</sup> Other investigators have provided evidence in support of the idea that Kir1.1 (the ROMK) is a pore-forming subunit of the  $mK_{ATP}$  channel.<sup>24</sup> In the current study, changes in mitochondrial and myocyte volume were observed in the presence of ROMK blocker TPN-Q in the presence of DZX. A dose effect on mitochondrial volume was not detectable using TPN-Q in the presence of DZX and multiple doses of TPN-Q. The stability of TPN-Q in the vehicle medium was thus evaluated in the media, in 20% acetonitrile, HEPES, and in water. However, no significant change in mitochondrial volume was observed after the addition of TPN-Q, even at high concentrations, in contrast to other investigators.<sup>24</sup> Thus, we conclude that TPN-Q had no effect on mitochondrial volume in the presence of DZX, suggesting a location of action of DZX that was distinct from the channels targeted by TPN-Q.

Myocytes demonstrated significant swelling in response to CPG that was prevented by the  $K_{ATP}$  channel opener, DZX, which is consistent with previous results.<sup>2,3</sup> Similarly, a correlation between myocyte volume changes and contractility changes was consistent with previous work, supporting this myocyte model of stunning.<sup>4,5</sup> TPN-Q prevented this beneficial



effect of DZX, thus implicating TPN-Q targets—Kir1.1, Kir3.1, or Kir3.4—in the cardioprotection of DZX at the cellular level. Other non-Kir TPN-Q targets or undefined channel subunits sensitive to TPN-Q may play a role (GIRK1/4,  $K_{ACH}$ , and voltage-dependent  $Ca^{2+}$ -activated  $K^{+}$  channels) in cardioprotection and will be the subject of future investigations.<sup>29–31</sup>

The present study further characterizes the mechanism of cardioprotection provided by DZX. TPN-Q did not alter DZX-induced mitochondrial swelling, but it did inhibit myocyte cardioprotection provided by DZX during stress. Given that TPN-Q inhibits Kir1.1-, Kir3.1-, and Kir3.4-dependent  $K^{+}$ -channel activities, these data support that any of these subunits (as well as undefined subunits sensitive to TPN-Q) could be involved in the cardioprotection afforded by DZX. However, these data also suggest that mitochondrial swelling by DZX does not involve Kir1.1, Kir3.1, or Kir3.4 and likely results from a yet to be identified mechanism.

Future work to identify the site and mechanism of action of  $K_{ATP}$  channel openers will involve indirect methods until genetic identification is accomplished.

## Sources of Funding

This study was supported by NIH RO1 HL098182-01A1 (Lawton), NIH 5T32HL007776 (Henn, Janjua), and the Barnes Jewish Hospital Foundation (Lawton).

## Disclosures

None.

## References

- Mizutani S, Prasad SM, Sellitto AD, Schuessler RB, Damiano RJ Jr, Lawton JS. Myocyte volume and function in response to osmotic stress: observations in the presence of an adenosine triphosphate-sensitive potassium channel opener. *Circulation*. 2005;112:1219–1223.
- Prasad SM, Al-Dadah AS, Byrd GD, Flagg TP, Gomes J, Damiano RJ Jr, Nichols CG, Lawton JS. Role of the sarcolemmal adenosine triphosphate-sensitive potassium channel in hyperkalemic cardioplegia-induced myocyte swelling and reduced contractility. *Ann Thorac Surg*. 2006;81:148–153.
- Mizutani S, Al-Dadah AS, Bloch JB, Prasad SM, Diodato MD, Schuessler RB, Damiano RJ Jr, Lawton JS. Hyperkalemic cardioplegia-induced myocyte swelling and contractile dysfunction: prevention by diazoxide. *Ann Thorac Surg*. 2006;81:154–159.
- Maffit SK, Sellitto AD, Al-Dadah AS, Schuessler RB, Damiano RJ Jr, Lawton JS. Diazoxide maintains human myocyte volume homeostasis during stress. *J Am Heart Assoc*. 2012;1:jah3-e000778 doi: 10.1161/JAHA.112.000778.
- Al-Dadah AS, Voeller RK, Schuessler RB, Damiano RJ Jr, Lawton JS. Maintenance of myocyte volume homeostasis during stress by diazoxide is cardioprotective. *Ann Thorac Surg*. 2007;84:857–862.
- Auchampach JA, Maruyama M, Caverio I, Gross GJ. Pharmacological evidence for a role of ATP-dependent potassium channels in myocardial stunning. *Circulation*. 1992;86:311–319.
- Galinanes M, Shattock MJ, Hearse DJ. Effects of potassium channel modulation during global ischaemia in isolated rat heart with and without cardioplegia. *Cardiovasc Res*. 1992;26:1063–1068.
- Grover GJ, Sleph PG, Parham CS. Nicorandil improves postischemic contractile function independently of direct myocardial effects. *J Cardiovasc Pharmacol*. 1990;15:698–705.
- Pignat J, Bourgoin J, Dumont L. Cold cardioplegia and the  $K^{+}$  channel modulator aprikalim (RP 52891): improved cardioprotection in isolated ischemic rabbit hearts. *Can J Physiol Pharmacol*. 1994;72:126–132.
- Shigematsu S, Sato T, Abe T, Saikawa T, Sakata T, Arita M. Pharmacological evidence for the persistent activation of ATP-sensitive  $K^{+}$  channels in early phase of reperfusion and its protective role against myocardial stunning. *Circulation*. 1995;92:2266–2275.
- Flagg TP, Enkvetchakul D, Koster JC, Nichols CG. Muscle  $K_{ATP}$  channels: recent insights to energy sensing and myoprotection. *Physiol Rev*. 2010;90:799–829.
- Shyng S, Nichols CG. Octameric stoichiometry of the  $K_{ATP}$  channel complex. *J Gen Physiol*. 1997;110:655–664.
- Nichols CG, Singh GK, Grange DK.  $K_{ATP}$  channels and cardiovascular disease: suddenly a syndrome. *Circ Res*. 2013;112:1059–1072.
- Flagg TP, Kurata HT, Masia R, Caputa G, Magnuson MA, Lefer DJ, Coetzee WA, Nichols CG. Differential structure of atrial and ventricular  $K_{ATP}$ : atrial  $K_{ATP}$  channels require SUR1. *Circ Res*. 2008;103:1458–1465.
- Anastacio MM, Kanter EM, Keith AD, Schuessler RB, Nichols CG, Lawton JS. Inhibition of succinate dehydrogenase by diazoxide is independent of the ATP-sensitive potassium channel subunit sulfonylurea type 1 receptor. *J Am Coll Surg*. 2013;216:1144–1149.
- Sellitto AD, Maffit SK, Al-Dadah AS, Zhang H, Schuessler RB, Nichols CG, Lawton JS. Diazoxide maintenance of myocyte volume and contractility during stress: evidence for a non-sarcolemmal  $K_{ATP}$  channel location. *J Thorac Cardiovasc Surg*. 2010;140:1153–1159.
- Sellitto AD, Al-Dadah AS, Schuessler RB, Nichols CG, Lawton JS. An open sarcolemmal adenosine triphosphate-sensitive potassium channel is necessary for detrimental myocyte swelling secondary to stress. *Circulation*. 2011;124:S70–S74.
- Anastacio MM, Kanter EM, Makepeace C, Keith AD, Zhang H, Schuessler RB, Nichols CG, Lawton JS. Cardioprotective mechanism of diazoxide involves the inhibition of succinate dehydrogenase. *Ann Thorac Surg*. 2013;95:2042–2050.
- Anastacio MM, Kanter EM, Makepeace CM, Keith AD, Zhang H, Schuessler RB, Nichols CG, Lawton JS. Relationship between mitochondrial matrix volume and cellular volume in response to stress and the role of ATP-sensitive potassium channel. *Circulation*. 2013;128:S130–S135.
- Hanley PJ, Mickel M, Loffler M, Brandt U, Daut J.  $K_{ATP}$  channel-independent targets of diazoxide and 5-hydroxydecanoate in the heart. *J Physiol*. 2002;542:735–741.
- Sato T, Sasaki N, Seharaseyon J, O'Rourke B, Marban E. Selective pharmacological agents implicate mitochondrial but not sarcolemmal  $K_{ATP}$  channels in ischemic cardioprotection. *Circulation*. 2000;101:2418–2423.
- Garlid KD, Paucek P, Yarov-Yarovsky V, Murray HN, Darbenzio RB, D'Alonzo AJ, Lodge NJ, Smith MA, Grover GJ. Cardioprotective effect of diazoxide and its interaction with mitochondrial ATP-sensitive  $K^{+}$  channels. Possible mechanism of cardioprotection. *Circ Res*. 1997;81:1072–1082.
- Garlid KD, Dos Santos P, Xie ZJ, Costa AD, Paucek P. Mitochondrial potassium transport: the role of the mitochondrial ATP-sensitive  $K^{+}$  channel in cardiac function and cardioprotection. *Biochim Biophys Acta*. 2003;1606:1–21.
- Foster DB, Ho AS, Rucker J, Garlid AO, Chen L, Sidor A, Garlid KD, O'Rourke B. Mitochondrial ROMK channel is a molecular component of mito $K_{ATP}$ . *Circ Res*. 2012;111:446–454.
- Halestrap AP, Clarke SJ, Khaliulin I. The role of mitochondria in protection of the heart by preconditioning. *Biochim Biophys Acta*. 2007;1767:1007–1031.
- National Research Council (US) Committee for the Update of Guide to Care and Use of Laboratory Animals. 8th ed. Washington, DC: The National Academy Press; 2011.
- Beavis AD, Brannan RD, Garlid KD. Swelling and contraction of the mitochondrial matrix. I. A structural interpretation of the relationship between light scattering and matrix volume. *J Biol Chem*. 1985;260:13424–13433.
- Drewnowska K, Clemons HF, Baumgarten CM. Prevention of myocardial intracellular edema induced by St. Thomas' Hospital cardioplegic solution. *J Mol Cell Cardiol*. 1991;23:1215–1221.
- Kitamura H, Yokoyama M, Akita H, Matsushita K, Kurachi Y, Yamada M. Tertiapin potently and selectively blocks muscarinic  $K^{+}$  channels in rabbit cardiac myocytes. *J Pharmacol Exp Ther*. 2000;293:196–205.
- Kanjhan R, Coulson EJ, Adams DJ, Bellingham MC. Tertiapin-Q blocks recombinant and native large conductance  $K^{+}$  channels in a use-dependent manner. *J Pharmacol Exp Ther*. 2005;314:1353–1361.
- Jin W, Klem AM, Lewis JH, Lu Z. Mechanisms of inward-rectifier  $K^{+}$  channel inhibition by tertiapin-Q. *Biochemistry*. 1999;38:14294–14301.

## **Adenosine Triphosphate–Sensitive Potassium Channel Kir Subunits Implicated in Cardioprotection by Diazoxide**

Matthew C. Henn, M. Burhan Janjua, Evelyn M. Kanter, Carol M. Makepeace, Richard B. Schuessler, Colin G. Nichols and Jennifer S. Lawton

*J Am Heart Assoc.* 2015;4:e002016; originally published August 24, 2015;

doi: 10.1161/JAHA.115.002016

The *Journal of the American Heart Association* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231  
Online ISSN: 2047-9980

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://jaha.ahajournals.org/content/4/8/e002016>

Subscriptions, Permissions, and Reprints: The *Journal of the American Heart Association* is an online only Open Access publication. Visit the Journal at <http://jaha.ahajournals.org> for more information.