

View this Article Submit to PLOS Get E-Mail Alerts Contact Us

<u>PLoS One</u>. 2017; 12(5): e0178538. Published online 2017 May 31. doi: <u>10.1371/journal.pone.0178538</u> PMCID: PMC5451061 PMID: <u>28562688</u>

Smooth muscle cell-specific *Notch1* haploinsufficiency restricts the progression of abdominal aortic aneurysm by modulating CTGF expression

Jaspreet Sachdeva,^{#1} Advitiya Mahajan,^{#2} Jeeyun Cheng,³ Jeremy T. Baeten,³ Brenda Lilly,³ Helena Kuivaniemi,⁴ and Chetan P. Hans^{1,2,3,¤*}

Michael Bader, Editor

¹Ohio State University, Columbus, Ohio, United States of America
²Cardiology, Medical Pharmacology & Physiology and Dalton Cardiovascular Research Center, University of Missouri, Columbia, Missouri, United States of America
³Center for Cardiovascular Research and The Heart Center, Nationwide Children's Hospital, Columbus, Ohio, United States of America
⁴Division of Molecular Biology and Human Genetics, Department of Biomedical Sciences, Faculty of Medicine and Health Sciences, Stellenbosch University, Tygerberg, South Africa
Max Delbruck Centrum fur Molekulare Medizin Berlin Buch, GERMANY
#Contributed equally.
Competing Interests: The authors have declared that no competing interests exist.
Conceptualization: CPH JS AM JC.
Data curation: CPH JS AM JC JTB BL HK.
Formal analysis: CPH JS BL HK.
Funding acquisition: CPH.
Investigation: CPH JS AM JC JTB.

Methodology: CPH JS AM JC JTB HK.

Project administration: CPH.

Resources: CPH BL HK.

Software: CPH JS AM JTB.

Supervision: CPH.

Validation: CPH JS AM JC JTB HK.

Visualization: CPH JS AM JC JTB HK.

Writing - original draft: CPH JS AM JC.

Writing - review & editing: CPH JS AM JC JTB BL HK.

^aCurrent address: School of Medicine-Cardiovascular Medicine, University of Missouri, Dalton Cardiovascular Research Center, Columbia, Missouri, United States of America

* E-mail: HansCP@health.missouri.edu

Received 2016 Oct 12; Accepted 2017 May 15.

Copyright © 2017 Sachdeva et al

This is an open access article distributed under the terms of the <u>Creative Commons Attribution License</u>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Aims

Infiltration of macrophages and apoptosis of vascular smooth muscle cells (VSMCs) promote the development of abdominal aortic aneurysm (AAA). Previously, we demonstrated that global Notch1 deficiency prevents the formation of AAA in a mouse model. Herein, we sought to explore the cell-specific roles of Notch1 in AAA development.

Methods and results

Cell-specific *Notch1* haploinsufficient mice, generated on *Apoe^{-/-}* background using Cre-lox technology, were infused with angiotensin II (1000 ng/min/kg) for 28 days. *Notch1* haploinsufficiency in myeloid cells (n = 9) prevented the formation of AAA attributed to decreased inflammation. Haploinsufficiency of *Notch1* in SMCs (n = 14) per se did not prevent AAA formation, but histoarchitectural traits of AAA including elastin degradation and aortic remodeling, were minimal in *SMC-Notch1^{+/-};Apoe^{-/-}* mice compared to *Apoe^{-/-}* mice (n = 33). Increased immunostaining of the contractile SMC-phenotype markers and concomitant decreased expression of synthetic SMC-phenotype markers were observed in the aortae of *SMC-Notch1^{+/-};Apoe^{-/-}* mice. Expression of connective tissue growth factor (CTGF), a matrix-associated protein that modulates the synthetic VSMC phenotype, increased in the abdominal aorta of *Apoe^{-/-}* mice and in the adventitial region of the abdominal aorta in human AAA. *Notch1* haploinsufficiency decreased the expression of Ctgf in the aorta and *in vitro* cell culture system. *In vitro* studies on SMCs using the Notch1 intracellular domain (NICD) plasmid, dominant negative mastermind-like (dnMAML), or specific siRNA suggest that Notch1, not Notch3, directly modulates the expression of CTGF.

Conclusions

Our data suggest that lack of Notch1 in SMCs limits dilation of the abdominal aorta by maintaining contractile SMC-phenotype and preventing matrix-remodeling.

Introduction

Abdominal aortic aneurysm (AAA) is a localized dilation of the abdominal aorta exceeding the normal diameter by more than 50% of its original size[$\underline{1}, \underline{2}$]. The prevalence of AAA ranges from 1.3% in men 45 to 54 years of age to ~9% in the population over 65 years of age[$\underline{2}, \underline{3}$]. In the United States, aortic aneurysms account for more than 10,000 deaths per year[$\underline{2}$]. Understanding the underlying mechanisms of AAA expansion will lead to novel pharmacologic therapies to reduce AAA development and rupture[$\underline{4}-\underline{6}$].

The development of AAA is characterized by transmural infiltration of macrophages and depletion of vascular smooth muscle cells (VSMCs) at the vascular injury site, which results in the focal enlargement of the abdominal aorta[7-10]. Excessive infiltration of macrophages and apoptosis of

VSMCs stimulates activation of matrix metalloproteinases (MMPs) resulting in the fragmentation of the elastin and interstitial collagenous matrix of the medial layer of the aorta[11]. VSMCs are capable of phenotypic modulation in response to different environmental cues. Contractile and synthetic VSMCs, which represent the two ends of a spectrum, have different morphological components and functional roles[12]. Contractile VSMCs are quiescent cells, display an elongated spindle-shaped morphology and express a unique repertoire of contractile proteins that serve as VSMC markers including SMA- α , SM22 α , smMHC and smoothelin[13, 14]. Vascular injury, results in abundant proliferative and migratory phenotypic modulation of VSMCs, which produce extracellular matrix (ECM) proteins such as elastin and collagen to provide tensile strength to the injured vessel wall. However, enhanced degradation of structural proteins, together with a reduced capacity to synthesize new matrix proteins, progressively weaken the aortic wall, resulting in rupture.

Notch1 signaling has gained substantial interest in the past few years because of its involvement in multiple aspects of vascular development and remodeling including myeloid and VSMC differentiation[15–23]. Notch1 signaling also regulates vascular inflammation[24], VSMC apoptosis[25, 26], and MMP-dependent proteolysis[27, 28]. Many of these pathways are implicated in the pathogenesis of aneurysms[29]. There are four types of Notch receptors (Notch1-4) expressed in all vascular cells including endothelial cells, VSMCs and macrophages[30]. Upon ligand receptor binding (Jagged or Delta-like), the Notch intracellular domain (NICD; active form of Notch1) is released from the membrane and is translocated into the nucleus where it binds to the transcriptional regulator RBPJ (also known as CBF-1). NICD activates the transcription of target genes by displacing the repressor complex from CBF-1 genes and recruiting activators including mastermind-like protein (MAML)[31]. Our laboratory identified Notch1 signaling as an important factor that mediates inflammatory response and disease initiation in angiotensin II (AngII) induced mouse model of AAA[32, 33]. We also reported that reconstitution of bone marrow-derived cells from Notch1+/-; Apoe-/- mice (donor) in irradiated Apoe^{-/-} mice (recipient) decreased the occurrence of aneurysm[<u>33</u>]. The diversified roles of Notch1 in various vascular cells demonstrate the complexity of Notch signaling and warrant studying the individual contributions of the Notch1 receptor in each vascular cell type on aneurysms.

The goal of this study was to define the unique contributions of SMC-specific and myeloid-specific Notch1 signaling in the development of AAA. We present evidence that myeloid-specific *Notch1* haploinsufficiency protects against the formation of AAA, whereas SMC-specific *Notch1* haploinsufficiency interferes with the progression of aortic aneurysms. Collectively, these findings imply that Notch1 signaling in different cell types affects the development of aortic aneurysm by multifarious mechanisms and represents a potential target to mitigate the deleterious effects of AAA at different stages of the disease progression.

Materials and methods

To determine the contributions of each vascular cell type in AAA development, we used a Cre-Lox recombination approach. The following cell-specific *Notch1* haploinsufficient mice were generated:

- 1. Myeloid specific Notch1 haploinsufficient mice (LysM-Cre).
- 2. Smooth muscle cell (SMC) specific Notch1 haploinsufficient mice (smMHC-Cre).
- 3. Endothelial cell (EC) specific Notch1 haploinsufficient mice (Tie2-Cre).

The *LysM-Cre*, *smMHC-Cre*, and *Tie2-Cre* mice were purchased from Jackson lab (Ann Arbor). Cell specific *Notch1* haploinsufficient mice were generated by crossing the offspring of *Notch1*^{flox/WT}; *Apoe^{-/-}* female mice with their respective *Cre* male mice against *Apoe^{-/-}* background as explained in <u>S1 Table</u>. All animals were genotyped by polymerase chain reaction (PCR) as shown in <u>S2</u>

and <u>S3</u> Tables). Aneurysmal studies were performed on these mice by infusing AngII or saline for 28 days using published protocols[<u>32</u>, <u>33</u>]. Briefly, mice were anesthetized in a closed chamber with 3% isoflurane in oxygen for 2 to 5 minutes until immobile. Each mouse was then removed, and taped on a heated (35–37°C) procedure board with 1.0–1.5% isoflurane administered via nosecone during minor surgery. After 28 days, mice were deep anesthetized with ketamine/xylazine (100 and 20 mg/kg, respectively) the aortae were dissected, fixed in 10% formalin, and maximal aortic diameters measured. The animal experiments were approved by Institutional Animal Care and Use Committee at the Research Institute at Nationwide Children's Hospital. All the animal experiments conform the NIH guidelines (Guide for the care and use of laboratory animals).

Isolation, culture, flow cytometry and co-culture of primary abdominal aorta-derived vascular smooth muscle cells (VSMCs) and bone marrow derived macrophages

Primary abdominal aorta-derived VSMCs were isolated from WT, Notch1^{+/-}, Apoe^{-/-} or Apoe^{-/-};Notch1^{+/-} mice (6–8 weeks of age). Briefly, mice were sacrificed with CO₂ asphyxiation and perfused with ice cold PBS through the left ventricle. Abdominal aorta was isolated and cleared from exogenous fat and adventitial layer followed by enzymatic digestion using collagenase/elastase (1 mg/mL collagenase, cat# LS004174, 0.744 units/mL elastase, cat# LS002279; Worthington) as described[34, 35]. VSMCs were grown in DMEM/F12 media without phenol red with 2% penicillin/streptomycin, 1% fungizone (Invitrogen), and 20% fetal bovine serum (Tissue Biologics) until cells were confluent, and then trypsinized (0.25%). Purity of SMC was confirmed by performing immunofluorescence on the isolated cells with antibodies to smooth muscle actin (SMA- α) as shown in our previous studies [35, 36]. All studies with VSMCs were performed on cells from passage 3–8. For apoptosis assay, VSMC were grown on 6 well plates. At \sim 70% confluence, the cells were treated with 1% FCS in DMEM containing lipopolysaccharide (LPS;100 ng/ml) for 24 hours. Afterwards, cells were stained with Annexin V Apoptosis Detection Kit (eBiosciences) using standard protocols. Flow cytometry was performed on a Becton Dickinson LSRII flow cytometer with DiVa software and data were analyzed with Flow Jo software (Ashland, OR) to quantify the percentage of differential apoptotic population. For co-culture assay, the bone marrow-derived macrophage (BMDMs) were isolated from *Notch1*^{+/-} and wild-type (*WT*) mice and differentiated to naïve macrophages onto transwell chambers using the standard protocol[32, 33]. The cells were then pretreated with vehicle or LPS (100 ng/ml) and interferon- γ (IFN- γ ; 20 ng/ml) for 3h, as described [37]. After pretreatment, the transwell chambers containing BMDMs were washed with media to remove excess of LPS/IFN- γ , and were transferred to another plate in which WT SMCs were previously grown at the ratio of 1:1. After 24 hours, the SMCs were washed and RNA was isolated for further experiments.

Transfection of smooth muscle cells

Plasmids and siRNA sequences were used for transfecting the cells. NICD and dnMAML plasmids were obtained from Dr. Lilly's laboratory. Predesigned Notch1 (s70698, Invitrogen) and Notch3 (s70707, Invitrogen) siRNA were used with Lipofectamine 3000 (L3000, Invitrogen) and Lipofectamine RNA iMAX (13778, Invitrogen). The cells were plated into 6-, 12-, or 24-well plates with their normal growth serum. Two hours prior to the transfection, DMEM media without serum and antibiotics were added to the cells. The reagents for transfection were prepared using OptiMEM media (31985–070, Invitrogen). Two different tubes were prepared for each well; one containing the OptiMEM and Lipofectamine P3000 and the other containing OptiMEM, Lipofectamine P3000/RNA iMAX reagent, and either the plasmid (2µl/µg DNA) or the siRNA. These two tubes were then combined and the mixture was incubated for 5 minutes at room temperature. This mixture was then added directly onto the cells containing the DMEM serum-free and antibiotic-free media. The media was changed four to six hours after transfection with normal growth medium.

Human aortic tissue samples

After written informed consent, full thickness infrarenal abdominal aortic wall tissue samples were collected from patients who underwent repair operations for AAA (n = 3, white men aged 67, 70, and 72 years) at the Harper University Hospital, Detroit, Michigan. Non-aneurysmal control samples were collected from the infrarenal segment of aorta (n = 3; white men aged 53, 53, and 78 years) at autopsies within 24 hours of death at the Detroit Coroner's Office. The samples were considered discarded tissue samples and no consent was required for the autopsy samples. At the time of harvesting, all samples were coded and no link to personal identifying information was kept. Tissue samples were incubated in phosphate-buffered formalin and embedded in paraffin for histological and immunohistochemical analyses. The collection of the tissue samples and their use for research projects was approved by the Institutional Review Board of Wayne State University, Detroit, Michigan, and the research carried out was in compliance with the Helsinki Declaration[32].

Histology, immunostaining, and quantitation

The mice from the experiments were euthanized and aortae were dissected. The thoracic and abdominal aortae were fixed in 10% formalin and serial sections were obtained. The abdominal aorta was cut into two equal halves and serial sections of the abdominal aorta ($5\mu M$) were prepared throughout the aneurysmal area of the abdominal aorta. Several sections of the abdominal aorta at regular intervals (200µM) were subjected to hematoxylin and eosin (H&E), elastin, and Masson's trichrome stain for histoarchitectural evaluation of aneurysm. The serial tissue sections obtained from these mice were further subjected to immunohistochemistry (IHC) or immunofluorescence (IF). For IHC, the abdominal aortae were stained with antibodies for active caspase-3, matrix metallopeptidase-2 (Mmp2), Mmp9, Mmp12, smooth muscle myosin heavy chain (smMHC), alpha-smooth muscle actin (SMA- α), osteopontin, vimentin, connective-tissue growth factor (Ctgf), NICD, interferon regulatory factor 8 (Irf8), Cd206, and monocyte chemotactic protein-1 (Mcp1). The intensity of the immunostaining was evaluated by obtaining 4–5 images from random areas of interest at 40X from each tissue (n = 4) and quantified using 'Fiji' version of Image J following the software directions[38]. For IF, CTGF and antifibroblast TE-7 were used to co-stain the sections of the human tissue. Antifade Mounting Medium containing DAPI (Vector Laboratories) was used to stain nuclei and mount the slides. The specificity of the antibody staining was determined using non-specific IgG against the source of host species. IHC and IF was performed using protocols described in a previous study[32, 33]. The list of antibodies can be found in <u>S4 Table</u>.

Protein isolation and Western blot

Total proteins were extracted from the cells with RIPA buffer supplemented with protease and phosphatase inhibitor cocktails (Roche). After homogenization, protein concentrations were measured and 20 μ g of protein were loaded onto 10% SDS-polyacrylamide gel, and transferred to PVDF membrane using standard protocols. The membranes were probed for CTGF and NICD using specific antibodies using standard protocol[<u>35–37</u>, <u>39</u>]. GAPDH antibody was used for internal loading control.

RNA isolation and qRT-PCR

Total RNA was isolated from the primary cell lines and the aortic tissue (suprarenal aorta approximately 5 mm thick) using TRIzol reagent (Ambion) according to manufacturer's protocol. The aortic tissue samples were first homogenized using TissueLyser II (Qiagen). Isolated RNA underwent a DNAse treatment using DNAse Treatment and Removal reagent (Ambion) to remove contaminating genomic DNA. cDNA was synthesized using the SuperScript VILO kit (Invitrogen) and subjected to quantitative PCR (qPCR) by SYBR Green (Applied Biosystems) using an Applied Biosystems 7500 Fast Real-Time

PCR System. The samples were run in triplicate and the fold-change was determined by normalizing CT values against *Rpl13a*. The primers are listed in <u>S5 Table</u>.

Vascular smooth muscle cell contractility assay

The ability of cells to contract a collagen gel matrix was determined using a 96-well collagen gel contraction assay using standard protocol [40]. Aortic VSMCs isolated from Apoe^{-/-} or Notch1^{+/-};Apoe^{-/-} mice were first cultured in a 6-well plate and treated with or without 10 ng/ml CTGF for 72 hours. Cells were then trypsinized, counted, and suspended in a collagen solution at a concentration of $6x10^5$ cells/ml. This collagen solution contained the following: 43% (v/v) sterile PBS (Hyclone), 34% (v/v) 3 mg/ml rat tail collagen (BD Biosciences), 13.6% (v/v) serum-free DMEM (Hyclone), 7.6% (v/v) 10x Medium 199 (GIBCO), and 1% (v/v) FBS (Mediatech). The solution was brought to neutral pH (\sim 7.0) by addition of NaOH using the phenol red indicator in the DMEM. Then, 70 µl of this collagen+cell solution was loaded into the wells of a 96-well plate and incubated at 37°C for 30 minutes to allow collagen to polymerize. A total of 140 µl of DMEM with 2% FBS was then added and the wells were incubated for 48 hours. The gels were then gently released from attachment to walls of the wells by Dumont #55 Forceps and 0 hour images were taken. The media was then replaced with 10% FBS DMEM to induce contraction and incubated for further 2 hours. After 2 hours, another image was taken of each well to determine the amount of gel contraction. Each treatment group was represented by 6 replicates in each experiment, and repeated in 3 independent experiments. The images were analyzed using ImageJ software[41] to measure the gel area at each time-point, from which the percentage decrease of gel area was determined [42].

Aortic ring assay

Aortae from *Apoe^{-/-}* male mice were isolated. These aortae were dissected to obtain 4–5 sections (rings) from the abdominal region for transfection. The aortic rings were treated with the following in the absence or presence of CTGF (Invitrogen): NICD and dnMAML plasmid (along with the empty vector), non-specific siRNA (4390846, Invitrogen), siRNA Notch1 (s70700, Invitrogen), and siRNA Notch3 (s70708, Invitrogen) using standard protocol[43]. Plasmids and pre-designed siRNA were transfected with Lipofectamine 3000 and Lipofectamine RNAiMAX, respectively. The rings were plated onto chamber slides in DMEM High Glucose media (Hyclone) with 10% FBS (Fisher Scientific). Two hours prior to the transfection, DMEM media without serum and antibiotics were added to the chamber slides. The reagents for transfection were prepared using OptiMEM media (Invitrogen). Two different tubes were prepared for each well; one containing the OptiMEM and Lipofectamine 3000 or RNA iMAX and the other containing OptiMEM, P3000 reagent, and either the plasmid or the siRNA. These two tubes were then combined and the mixture was incubated for 5 minutes at room temperature. This mixture was the added directly onto the chamber slides with the aortic rings containing the DMEM serum-free and antibiotic-free media. The media was changed 24 hours after transfection to DMEM High Glucose media with 10% FBS. The aortic rings were then treated using recombinant human CTGF protein (PHG0286; Gibco) for additional 48 hours. The rings were then collected and processed for histology.

Statistical analysis

Statistical comparisons were performed using either the Student t-test or 1-way ANOVA followed by the Bonferroni multiple comparison correction. GraphPad Prism version 5.0 (GraphPad Software, Inc., CA) was used for these comparisons and P<0.05 was considered significant. Assumptions of normality and equal variance were tested and satisfied using SAS software version 9.3 (SAS Institute Inc., NC). For multiple comparisons, a post-Bonferroni test with Holm correction (for 3 comparisons) was

performed for maximal aortic width graph and survival graph. For the real-time RT-PCR quantification, we performed the Kruskal–Wallis test using a non-parametric method for the overall significance of the data and ordinary ANOVA followed by a Bonferroni-Holm multiple comparisons test with single-pooled variance for multiple comparisons using GraphPad Prism version 5.0. For the statistical analysis of actual incidence, Fisher's exact test was used with the SAS software. The actual incidence of the disease is defined as 50% of greater increase in the maximal aortic width compared with the control group.

Results

Cell-specific *Notch1* haploinsufficiency differentially interferes with AAA development in AngII-induced mouse model

To determine the contributions of SMCs, ECs, and macrophages in AAA development, we used the Cre-Lox recombination approach. The macroscopic quantification of the external diameter of suprarenal aorta demonstrated a significant exacerbation in maximal aortic width of Apoe^{-/-} mice with AngII treatment to 2.37 ± 0.58 mm (n = 24, P<0.001; Fig 1B, 1E and 1F) compared to saline treated Apoe^{-/-} mice $(1.10 \pm 0.10 \text{ mm}; n = 6; \text{Fig.} 1\text{A})$. No difference in the littermate controls was observed in the Apoe^{-/-} mice for each Cre or Notch l^{flox/wt} with regard to aneurysm, hence all these mice are pooled in one group (n = 33). The maximal aortic width in *myeloid-Notch1*^{+/-}; Apoe^{-/-} mice remained significantly lower as compared to AngII-treated Apoe^{-/-} mice and was comparable to saline-infused Apoe^{-/-} mice $(0.97 \pm 0.18; n = 7, P < 0.001, Fig 1D and 1F)$. Interestingly, in SMC-Notch1^{+/-}; Apoe^{-/-} mice, dilation of aorta increased moderately and remained in a narrow range. At day 28 the maximal aortic width in *SMC-Notch1^{+/-};Apoe^{-/-}* mice was modestly but insignificantly lower than AngII-treated *Apoe^{-/-}* mice $(1.83 \pm 0.47; n = 7, Fig 1C, 1E and 1F)$. The maximal aortic width in EC-Notch1^{+/-}; Apoe^{-/-} mice was comparable to AngII-treated Apoe^{-/-} mice (S1 Fig; n = 7). Overall, AngII treatment resulted in 27% mortality in Apoe^{-/-} mice within 10 days of the treatment (Fig 1G). In SMC-Notch1^{+/-}; Apoe^{-/-} mice, 50% mortality was observed and occurred within 7 days of treatment, but was not statistically significant as compared to Apoe^{-/-} mice with AngII treatment (Fig 1G), which may be attributed to either initial influx of Notch1 sufficient macrophages at the vascular lesion or some unrelated effects of *Notch1*-haploinsufficiency[44]. In our studies, the aortae from dead SMC-Notch1^{+/-}; Apoe^{-/-} mice had dissection in the thoracic region. Indeed, various studies have shown strong correlation between Notch1 mutation and thoracic aortic aneurysm (TAA) suggesting that TAA and AAA represent a very complex pathology with regarding to Notch1 signaling pathway[45-47]. Interestingly, the dilation of proximal aorta in the ascending aorta and arch was comparable in Apoe^{-/-} and SMC-Notch1^{+/-}; Apoe^{-/-} mice at day 28 of AngII treatment and was larger than in *myeloid-Notch1^{+/-};Apoe^{-/-}* mice (S2 Fig). The trend in the mortality in EC-Notch1^{+/-}; Apoe^{-/-} mice was similar to SMC-Notch1^{+/-}; Apoe^{-/-} mice (S1 Fig). The mortality in *myeloid-Notch1*^{+/-}; Apoe^{-/-} mice was comparable to Apoe^{-/-} mice in response to AngII (33%), whereas no deaths were observed in the saline-treated mice (Fig 1G). Summarizing these findings, Notch1 haploinsufficiency in macrophages is sufficient to prevent the formation of AAA in Apoe^{-/-} mice in response to AngII, whereas Notch1 haploinsufficiency in SMCs may interefere with the progression of AAA.



<u>Fig 1</u>

Open in a separate window

Cell-specific *Notch1* haploinsufficiency differentially interferes with AAA development in AngII-induced mouse model.

Representative aortae isolated from *Apoe^{-/-}* mice treated with saline (**A**), AngII (**B**), *SMC-Notch1^{+/-};Apoe^{-/-}* mice (**C**;) and *myeloid-Notch1^{+/-};Apoe^{-/-}* mice (**D**;) treated with AngII (28 days). Images were taken using Zeiss Stemi 2000-C microscope. **Scale bar, 1 mm**. (**E-F**), Quantitative measurement of maximal aortic width (mm) in different groups. Each symbol represents an individual animal. Mean and SEM are shown. (G) Survival graph showing the mortality of individual mice in each group during 28 day AngII infusion. ***P < 0.001 (post-Bonferroni test with Holm correction).

Cell-specific *Notch1* haploinsufficiency prevents elastin degradation and limits matrix metalloproteinase production

The mouse experimental tissues were examined histologically for the characteristic architecture and distinct features of aneurysm. HE staining of the abdominal aorta after 28 days of AngII infusion in Apoe^{-/-} mice showed dilation of the aortic wall, inflammatory cell infiltration, and disarrangement of ECM with clear signs of elastin fragmentation (Fig 2B, 2F and 2J) as compared to saline-infused Apoe^{-/-} mice (Fig 2A, 2E and 2I). No such histoarchitectural features of aneurysm were observed in the aortae of myeloid-Notch1^{+/-}; Apoe^{-/-} mice infused with AngII (Fig 2D, 2H and 2L) which displayed a well-defined lumen, no visible elastin fragmentation and minimal inflammatory cell infiltration. SMC-Notch $I^{+/-}$; Apoe^{-/-} mice depicted a distinct morphology of abdominal aorta associated with moderate adventitial thickening, marginal aortic remodeling and no visible signs of elastin fragmentation (Fig 2C, 2G and 2K). Moderate to high collagen content was observed in the adventitial region of abdominal aorta in AngII-treated Apoe^{-/-} mice (Fig 2N) as compared to Apoe^{-/-} mice treated with saline (Fig 2M). Collagen content in the adventitial aorta of SMC-Notch1^{+/-}; Apoe^{-/-} mice were comparable to AngII-treated Apoe^{-/-} mice (Fig 20). However, the mRNA expression Colla1, Colla2, and Col3a1 increased disproportionately by almost 4-fold in response to AngII in Apoe^{-/-} mice compared with AngII-treated SMC-Notch1^{+/-}; Apoe^{-/-} mice, suggesting increased collagen degradation and turnover in the former group (Fig 2Q-2S). Collagen content was not considerably altered in the aortae of mveloid-*Notch1*^{+/-};*Apoe*^{-/-} mice infused with AngII (Fig 2P–2S).



Open in a separate window

<u>Fig 2</u>

Cell-specific *Notch1* haploinsufficiency in vascular smooth muscle cells and macrophages protects against extensive structural damage and collagen deposition in the abdominal aortic wall in response to AngII (28 days) treatment.

(A-D) Cross-section of representative aortae stained with hematoxylin and eosin (HE) staining showing lumen and intraluminal thrombus. Panel (E-H) Representative photomicrographs showing transmural inflammatory cell infiltration. (I-L) Representative images of modified Verhoeff Van Gieson staining demonstrating the extent of elastin fragmentation; (M-P) Collagen contents in the abdominal aorta of experimental groups visualized in blue using trichrome staining. Scale bar = 50 µm in A-P. (Q-S) Bar graphs represent gene expression of *Col1a1*, *Col1a2* and *Col3a1* in the aortae of the *Apoe^{-/-}* and cell-specific *Notch1* haploinsufficient *Apoe^{-/-}* mice after 28 days AngII treatment. The results were standardized

to *Rpl13a* and reported as ratio (mean \pm SEM, n = 3 for each group) to saline-treated mice. ****P*<0.001 (ordinary ANOVA followed by a Bonferroni-Holm multiple comparisons test).

MMPs have been implicated in degrading a wide range of ECM proteins in vascular diseases including AAA. As assessed by the IHC, a high immunoreactivity of Mmp2 and Mmp9 was observed throughout the medial and adventitial layers of abdominal aorta of $Apoe^{-/-}$ mice with AngII treatment (S3B and S3F Fig) as compared to saline treated $Apoe^{-/-}$ mice (S3A and S3E Fig). In contrast, minimal Mmp2 and Mmp9 immunoreactivity was present in the abdominal aortae of *SMC-Notch1*^{+/-};*Apoe*^{-/-} and *myeloid-Notch1*^{+/-};*Apoe*^{-/-} mice in response to AngII (S3C, S3D, S3G and S3H Fig).

Apoptotic cell death and depletion of medial SMC has been shown to play an important role in human AAA pathological samples and in mouse aneurysmal models[48, 49]. Immunohistochemical analyses showed increased active caspase-3 immunoreactivity in the medial layer of aorta in the $Apoe^{-/-}$ mice infused with AngII compared with saline treated $Apoe^{-/-}$ mice (S3I and S3J Fig). Minimal active caspase-3 immunostaining was observed in the abdominal aorta of SMC-Notch1^{+/-}; $Apoe^{-/-}$ and myeloid-Notch1^{+/-}; $Apoe^{-/-}$ mice (S3K and S3L Fig). Apoptotic cell death of LPS-stimulated VSMC isolated from the abdominal aorta of $Notch1^{+/-}$; $Apoe^{-/-}$ mice as detected by FACS analysis using annexin V-propidium iodide (PI) staining (S4A–S4C Fig). These data suggest that SMC- and myeloid-specific Notch1 haploinsufficiency impact the pathological sequelae associated with AAA by different mechanisms. In following experiments, we focused on the mechanism by which SMC-specific Notch1 haploinsufficiency interferes with the development of AAA.

Shift in SMC phenotype in SMC-specific *Notch1* haploinsufficient mice is associated with decrease in CTGF expression

Nuclear Notch1 staining representing active NICD was observed in the aortae of Apoe^{-/-} mice (arrows in S5B Fig). The medial layers of aortae in SMC-Notch $1^{+/-}$; Apoe^{-/-} mice were devoid of such nuclear Notch1 staining (arrows in S5C Fig). Surprisingly, no nuclear NICD staining was observed in the medial layer of *myeloid-Notch1*^{+/-}; Apoe^{-/-} mice in spite of cytoplasmic presence of NICD (arrows in S5D Fig). The results confirm that depletion of Notch1 signaling reduces the expression of Notch1 activity in respective cell-specific Notch1 haploinsufficient mice. We previously reported that Notch1 deficiency protects against AAA development by reducing M1 polarization of $M\phi$ [32, 33]. Increased expression of Irf8 and Mcp1 was observed in the abdominal aortae of Apoe^{-/-} mice in response to AngII (S5E-S5L Fig). The expression of these cytokines was minimal in the aorta of myeloid-Notch1^{+/-};Apoe^{-/-} mice, with concomitant increase in the expression of Cd206. SMC-Notch1^{+/-};Apoe^{-/-} mice depicted a distinct phenotype with regard to the expression of macrophage polarization. Whereas, the expression of Irf8 and Mcp1 was comparable to Apoe^{-/-} mice in response to AngII, the expression of Cd206 was also increased in the adventitia suggesting heterogeneity in the macrophage population (S5G, S5K and S5O Fig). The gene expression of iNos (a M1 macrophage marker) was significantly increased in the Apoe^{-/-} mice in response to AngII compared to other groups, whereas in the myeloid-Notch1^{+/-};Apoe^{-/-} mice, expression of Arg1 (a M2 macrophage marker) was significantly increased compared to the other groups (S5Q and S5R Fig).

Next, we examined the factors that define a contractile or synthetic SMC phenotype. Expression of the contractile markers smMHC and SMA- α were highly expressed in the aortae of control *Apoe^{-/-}* mice (<u>Fig 3A</u> and <u>S6A Fig</u>). Decreased immunoreactivity for smMHC and SMA- α was observed in the medial layer of abdominal aorta in response to AngII in *Apoe^{-/-}* mice (<u>Fig 3B and 3E</u> and <u>S6B Fig</u>). *Notch1* haploinsufficiency in SMCs prevented such loss in the contractile SMC-phenotype markers (Fig 3C, and S6C Fig). Remarkably, the adventitial thickening observed in these groups was also associated with increased expression of smMHC and SMA- α , and was completely absent in the *Apoe*^{-/-} mice in response to AngII (red arrows in Fig 3C and 3E and S6C Fig). Expression of contractile markers in the *myeloid-Notch1*^{+/-};*Apoe*^{-/-} mice was similar to *SMC-Notch1*^{+/-};*Apoe*^{-/-} mice (Fig 3D and 3E and S6D Fig). However no migration of these contractile SMCs was observed in the adventitial of *myeloid-Notch1*^{+/-};*Apoe*^{-/-} mice. Conversely, the expression of osteopontin and vimentin, markers of a synthetic-SMC phenotype was increased with AngII infusion primarily in the adventitial region in the aorta of *Apoe*^{-/-} mice (Fig 3F and 3J and S6F Fig). The expression of these synthetic-SMC markers was minimal in the abdominal aorta of *SMC-Notch1*^{+/-};*Apoe*^{-/-} mice. In the *myeloid-Notch1*^{+/-};*Apoe*^{-/-} mice, the expression of osteopontin appears to be similar to *Apoe*^{-/-} mice with AngII infusion. Mmp12 plays an important role in SMC migration and proliferation during vascular remodeling. As shown in Fig 3K–3O, AngII caused induction of Mmp12 expression in both medial and adventitial layers of abdominal aorta in the *Apoe*^{-/-} mice compared to control mice. No such increase in Mmp12 expression was observed in the abdominal aorta of *SMC-Notch1*^{+/-};*Apoe*^{-/-} mice in response to AngII (Fig 3M).



Open in a separate window

<u>Fig 3</u>

SMC-specific Notch1 haploinsufficiency dependent regulation of SMC phenotype is associated with changes in CTGF expression.

Representative images of IHC staining of abdominal aorta from respective experimental groups with antibodies against smMHC (**A-D**) and osteopontin (**F-I**) as markers of contractile or synthetic-phenotype marker of SMCs, respectively. IHC staining with antibody against Mmp12 (**K-N**), and Ctgf (**P-S**) performed on cross sections of abdominal aortae isolated from $Apoe^{-/-}$ cell-specific *Notch1* haploinsufficient $Apoe^{-/-}$ mice after 28 days AngII treatment. **Scale bars = 50** µm. The IHC for respective immunostaining was quantified using 'Fiji' version of image J (**E, J, O, T**). Bar graphs represent fold

change in gene expression of $Ctgf(\mathbf{W})$, $Tgf-\beta 1(\mathbf{X})$, $Tgf-\beta 2(\mathbf{Y})$, $Smad2(\mathbf{Z})$ and $Smad3(\mathbf{AA})$ using mRNA obtained from aortae of $Apoe^{-/-}$ and cell-specific *Notch1* haploinsufficient mice. The results were standardized to Rpl13a and reported as ratio to saline-treated mice (mean \pm SEM, n = 3 for each group). (**AB**) Fold change in Ctgf expression in SMCs isolated from Apoe^{-/-} and *Notch1*^{+/-}; $Apoe^{-/-}$ mice at basal levels and in the presence of TNF- α . ***P<0.001; **P<0.01; and *P<0.05 (ordinary ANOVA followed by a Bonferroni-Holm multiple comparisons test).

Connective tissue growth factor (CTGF), secreted by synthetic VSMCs and fibroblasts, is a potent ECM-inducing growth factor involved in regulating aneurysm-associated vascular remodeling [50, 51]. We examined if *Notch1* haploinsufficiency in SMCs is associated with alterations in Ctgf expression. Minimal immunostaining of Ctgf was observed in the saline infused Apoe^{-/-} mice (Fig 3P). In response to AngII, Ctgf immunoreactivity markedly increased in the abdominal aorta of Apoe^{-/-} mice (Fig.3T) and the distribution was similar to osteopontin and vimentin expression. Moderate expression of Ctgf was also observed in the aorta of *mveloid-Notch1*^{+/-}; Apoe^{-/-} mice with AngII treatment (Fig 3Q and 3T). Immunoreactivity of Ctgf was minimal in the aorta of SMC-Notch1^{+/-};Apoe^{-/-} mice, suggesting a phenotypic shift with SMC-specific Notch1 haploinsufficiency (Fig 3R). Next, we examined if Notch1 deficiency causes any alterations in the gene expression of canonical Tgf- β signaling, an upstream regulator of Ctgf pathway. Gene expression of Ctgf, Opn, Mmp12, Tgf- $\beta 1$ and Tgf- $\beta 2$ was increased in the aorta of *Apoe^{-/-}* mice with AngII infusion (Fig.3U-3Y). Myeloid-specific *Notch1* haploinsufficiency did not prevent such an increase in the gene expression of Ctgf and Tgf- βI . In fact, the expression of *Tgf-* β 2 was even higher in *myeloid-Notch* $I^{+/-}$; *Apoe*^{-/-} mice than *Apoe*^{-/-} mice with AngII. Conversely, the expression of Ctgf, Tgf- $\beta 1$ and Tgf- $\beta 2$ was significantly decreased in the aortae of SMC-Notch1^{+/-};Apoe^{-/-} mice as compared to Apoe^{-/-} mice (Fig.3W-3Y). Expression of Smad2, a downstream target of Tgf- β 2 signaling was also increased in *Apoe^{-/-}* mice and *myeloid-Notch1^{+/-};Apoe^{-/-}* mice with AngII treatment and was significantly reduced in SMC-Notch1^{+/-}; Apoe^{-/-} mice (Fig 3Z). Surprisingly, expression of Smad3 was significantly increased in SMC-Notch1^{+/-}; Apoe^{-/-} mice (Fig 3AA). The gene expression of Ctgf was also lower in the SMCs isolated from the abdominal aorta of Notch1^{+/-}; Apoe^{-/-} mice as compared to Appe^{-/-} mice at basal conditions or in the presence of TNF- α (10 ng/ml for 24 hours; Fig 3A and 3B).

CTGF has been shown to be a direct modulator of the VSMC phenotype in vascular diseases in human[50]. A significant increase in CTGF-expressing cells was observed in AAA in the inflammatory regions compared with non-inflammatory regions in the same AAA tissue or in the aorta of control human subjects (Fig 4A-4D). The inflammatory and non-inflammatory regions in these tissues were recognized by the presence of macrophages and other inflammatory cells[32]. Next, we examined if increased CTGF expression is associated with synthetic SMC-phenotype in the aorta of human AAA patients. Co-immunostaining with a highly specific TE-7 antibody which identifies fibroblast-like synthetic SMCs[52] suggests that CTGF expression is highly correlated with fibroblast-like synthetic SMC-phenotype (Fig 4E-4P). To further examine if macrophages modulate SMC-phenotype, we performed co-culture assay. The SMCs were obtained from the abdominal aorta of wild type (WT) mice and bone marrow derived macrophages (BMDMs) were isolated from either WT or $Notch1^{+/-}$ mice. The BMDMs were pretreated with LPS/IFN-y to polarize them into pro-inflammatory M1 phenotype as explained [32, 33]. As the data demonstrate, naïve BMDMs from either WT or Notch $1^{+/-}$ mice did not affect the expression of smMHC, Ctgf, Mmp12, or vimentin in SMCs. Co-culture with BMDMs, pretreated with LPS/IFN- γ showed no significant changes in the expression of *Mmp12* or *vimentin*. The expression Ctgf and smMHC, however was decreased in the WT SMCs in response to co-culture with *Notch1^{+/-}* BMDM pretreated with LPS/IFN- γ . These data support the hypothesis that Notch1 deficiency

may be directly regulating SMC phenotype and that these effects may be mediated independent of its effects on macrophage polarization.



<u>Fig 4</u>

CTGF expression in human AAA correlates strongly with synthetic SMC-phenotype.

(A-D) Immunohistochemical staining for CTGF on the normal and aneurysmal human abdominal aorta sections. (E-P) Micrographs showing immunofluorescence staining for CTGF (green; E, I, M) and fibroblast marker TE-7 (red; E, J, N) on the tissue sections obtained from medial layer of healthy human controls (E-H) or aneurysmal (I-L) or non-aneurysmal aorta (M-P) from same human AAA. Merged images of CTGF and TE-7 staining from respective tissues (yellow; G, K, O) and nuclear staining (DAPI; H, L, P) are shown. Scale bars = 50 μm.

Gain- and loss-of-function studies suggest direct involvement of Notch1 in regulating CTGF expression

TGF- β 2 is a potent stimulator of SMC phenotypic modulation and is also upstream of CTGF signaling. We determined if Notch1 is upstream regulator in the Tgf- β 2 mediated regulation of CTGF. First, we confirmed the expression of CTGF in cultured human aortic SMCs (hASMCs) and mouse primary fibroblasts at basal levels and in response Tgf- β 2 agonist or antagonist. At basal conditions, very low expression of CTGF was observed and DAPT (a potent Notch inhibitor) further reduced CTGF expression marginally in these cell types (Fig 5A and 5B). In response to TGF- β 2 agonist (human recombinant protein), the CTGF expression increased about 2-fold in hASMCs and 100-fold in mouse

fibroblasts. Pre-treatment of hASMCs with DAPT significantly reduced such induction of CTGF expression (Fig 5A and 5B). In contrast, SB431542, an inhibitor of activin receptor-like kinase (ALK5; the TGF- β type I receptor) reduced the expression of CTGF in both cell types ~50% below basal expression. No significant effect of DAPT was observed in response to SB431542 probably because of very low expression of CTGF.



<u>Fig 5</u>

Gain- and loss-of-function studies suggest direct involvement of Notch1 in regulating CTGF expression.

(A-B) Bar graphs represent fold change in gene expression of CTGF in the human aortic smooth muscle cells (A) and mouse embryonic fibroblast (B) treated with DAPT for 24 hours followed by TGF- β agonist or inhibitor SB431542 [activin receptor-like kinase (ALK5; the TGF-beta type I receptor inhibitor]

treatment for 24 hours. **(C-F)** Graphs represent fold change in the expression of Notch1, CTGF, Notch3 and smoothelin in the human aortic smooth muscle cells 48 hours post transfected with Notch1 *intracellular domain* (NICD) overexpressing plasmid or dominant negative mastermind like proteins (dnMAML). **(G-J)** Graphs represent fold change in the expression of Notch1, CTGF, Notch3 and smoothelin in the human aortic smooth muscle cells 48 hours post transfected siRNAs for Notch1/Notch3.The qPCR data were standardized to *RPL13a* and reported as ratio (mean \pm SEM, n = 3 for each group) to empty plasmids for NICD/dnMAML or non-specific siRNA. ****P*<0.001, ***P*<0.01, **P*<0.05 (ordinary ANOVA followed by a Bonferroni-Holm multiple comparisons test). (EP = empty plasmid).

Next, we performed transfection studies on hASMCs using plasmids for NICD or a dominant negative form of the MAML activator (dnMAML) to overexpress or downregulate Notch signaling respectively. NICD-plasmid increased the expression of Notch1 by almost 250-fold and CTGF expression by 2-fold in the hASMCs (Fig 5C and 5D). Conversely, dnMAML-mediated knockdown of Notch signaling did not significantly affect CTGF expression (Fig 5C and 5D). In contrast, siRNA-mediated specific knockdown of Notch1 resulted in a more than 50% reduction in CTGF expression (Fig 5G and 5H). The discrepancy between the dnMAML and siRNA for Notch1 may be attributed to either inhibitory effects of dnMAML on other Notch family members including Notch2 and Notch3, or potential noncanonical activities of Notch1 that are affected by dnMAML. Further, siRNA for Notch3 did not affect the expression of CTGF significantly (Fig 51). However, decreased expression of smoothelin (a contractile smooth muscle marker) was observed with Notch3 siRNA (Fig 5J). Interestingly, the expression of Notch3 and smoothelin was also increased by almost 2-fold with NICD overexpression (Fig 5E and 5F). No synergistic effects of Notch1 and Notch3 inhibition were observed on the expression of CTGF and smoothelin as determined by the treatment of hASMCs with a combination of siRNA for both Notch1 and Notch3 (Fige 5G–5J). Next, we investigated if the alteration in gene expression of CTGF is translated in the change in protein content. Western Blotting was performed to observe this change. Reduced protein contents of CTGF were observed in Notch1^{+/-} SMCs and hASMCs treated with DAPT (S8 Fig). In accordance with the gene expression data, Tgf-B2 increased the CTGF protein contents, whereas reduced CTGF contents were observed in hASMCs with SB-431542, an inhibitor of Tgf- β signaling (S8A and S8B Fig). Reduced CTGF content was observed in SMCs with Notch1 deficiency under all these conditions. Overexpression of NICD reversed the protein contents of CTGF in hASMCs (S8C Fig). Collectively, our data suggest that Notch1 may be directly involved in the CTGF-associated hASMC phenotype modulation.

Notch1 deficiency reduces the CTGF-induced collagen gel stiffness in VSMCs

Previous studies have demonstrated that Notch1 signaling directly regulates CTGF in osteoclasts which increases fibrotic response[53]. To determine whether CTGF functionally modulates a shift towards a synthetic phenotype, we performed collagen gel contraction assay using VSMCs from the abdominal aortae of $Apoe^{-/-}$ and $Notch1^{+/-};Apoe^{-/-}$ mice. At basal condition, the contraction in the collagen of the VSMCs from $Apoe^{-/-}$ mice was ~13% in response to 10% FBS (Fig 6A and 6B) whereas the contraction in the collagen of the VSMCs from $Notch1^{+/-};Apoe^{-/-}$ mice was significantly higher at ~35% (Fig 6E and 6F; P<0.001). The contraction of collagen gel was reduced to just ~5% in the presence of CTGF in $Apoe^{-/-}$ VSMCs (Fig 6C and 6D), whereas in the VSMCs from $Notch1^{+/-};Apoe^{-/-}$ mice, this contraction was significantly higher at ~10% (Fig 6G and 6H; P<0.05). We further characterized the inhibitory effect of Notch1 on CTGF-induced changes in VSMC-phenotype using *ex vivo* aortic ring culture from $Apoe^{-/-}$ mice and NICD plasmid or siRNA constructs[43]. Overexpression of Notch1 increased the collagen content in the adventitial area of aortic rings as compared to the non-NICD

transfected aortic rings (marked with red boundary) at basal conditions (<u>Fig 6K</u>) and in the presence of human recombinant CTGF (<u>Fig 6O</u>). Transfection with siRNA Notch1 abrogated such increase in the collagen content at basal levels (<u>Fig 6L</u>) and in response to recombinant CTGF (<u>Fig 6P</u>) suggesting inhibitory effects of Notch1 deficiency on collagen content. Modest increase in the adventitial collagen content was observed in the aortic rings treated with siRNA for Notch3 and CTGF (<u>Fig 6Q</u>) suggesting differential effects of Notch isoforms. No difference was observed in the collagen content in the aortic rings treated with empty plasmid for NICD or non-specific siRNA controls (data not shown).



<u>Fig 6</u>

Notch1 deficiency reduces the CTGF-induced collagen gel stiffness in SMCs.

(A-H) Photographs depict contractility of collagen gel embedded with VSMCs isolated from $Apoe^{-/2}$ and Notch1^{+/-}; Apoe^{-/-} mice. VSMCs cultured in a 6-well plate and treated with or without CTGF (10 ng/ml) for 72 hours and resuspended in a collagen solution at a concentration of 6×10^5 cells/ml for 48 hours. After imaging, the gels were gently released from attachment and the media was replaced with 10% FBS DMEM to induce contraction and incubated for 2 hours. The gel was imaged to determine the amount of gel contraction. Each group is represented by 6 replicates in each experiment, and repeated in 3 independent experiments. The images were analyzed using ImageJ software to measure the gel area at each time-point, from which the percentage decrease of gel area was determined. The yellow dotted lines represent the area of the collagen gel. Bar graphs represent percent decrease of gel area in different treatment groups. (J-Q) Trichrome staining of the aortic rings from Apoe^{-/-} mice transfected with NICD, siRNA for Notch1 or Notch3 in the absence (J-M) or presence of CTGF (N-Q). The aortic rings were transfected with NICD, non-specific siRNA (4390846, Invitrogen), siRNA Notch1 or siRNA Notch3 for 24 hours. The aortic rings were then treated using recombinant human CTGF protein for additional 48 hours. The rings were then collected and processed for histology. The red dotted lines represent the adventitial layer enriched with collagen contents. Scale bars = 50 μ m. ***P<0.001, *P<0.05 (ordinary ANOVA followed by a Bonferroni-Holm multiple comparisons test).

These findings, coupled with our observations that Notch1 overexpression is correlated with CTGF upregulation, led us to propose that Notch1 signaling is a direct modulator of VSMC-phenotype. These data demonstrate the ability of CTGF to inhibit the contractility behavior of VSMCs and that inhibition of Notch1 signaling prevents such effect in a TGF- β signaling-dependent manner, which may play a role in maintaining VSMCs in contractile phenotype.

Discussion

In the current study, we demonstrated that *Notch1* haploinsufficiency in myeloid cells (LysM-Cre) prevented the formation of AAA and recapitulated histological traits of global Notch1 deficiency including increased M2-differentiation of macrophages and decreased inflammation. Haploinsufficiency of Notch1 in SMCs (smMHC-Cre) did not prevent the AAA formation in these mice. However, typical histological characteristics of AAA including elastin fragmentation and aortic remodeling were minimal in SMC-Notch1^{+/-}; Apoe^{-/-} mice as compared to Apoe^{-/-} mice with AngII. Also, Notch1 haploinsufficiency in SMCs was associated with increased expression of contractile markers of SMCphenotype and concomitant decrease in the expression of markers of synthetic SMC-phenotype in the aorta. Expression of CTGF, an ECM-associated protein that modulates synthetic VSMC phenotype, was significantly increased in the abdominal aorta of Apoe^{-/-} mice in response to AngII. SMC-specific Notch1 haploinsufficiency prevented such increase in CTGF expression. Increased expression of CTGF was also observed in the adventitial region of the abdominal aorta in human AAA and correlated strongly with synthetic SMC-phenotype. In vitro studies on hASMCs and fibroblasts suggest that Notch1 modulates the expression of CTGF by a TGF- β dependent mechanism, which seems to be independent of interactions between SMCs and macrophages. Our data suggest that lack of Notch1 in myeloid cells is sufficient to prevent the formation of AAA whereas SMC-specific Notch1 haploinsufficiency limits dilation of the abdominal aorta by maintaining contractile SMC phenotype and preventing matrix remodeling (Fig 7).



The dashed arrows depict the mechanistic pathways to be tested in the study. We propose that myeloid specific *Notch1* haploinsufficiency prevents aneurysm development by modulating macrophage polarization. SMC-specific *Notch1* haploinsufficiency interferes with the progression of aneurysm by modulating CTGF-dependent SMC-phenotype. (Mφ: macrophages, SMC: smooth muscle cells; CTGF, connective tissue growth factor, opn: osteopontin,vim: vimentin).

Extensive studies using mouse models of AAA and our published data using bone marrow transplantation approaches have shown widespread inflammatory macrophages as major cellular components at the onset of the disease [10, 32, 33, 54]. Progression of aneurysm is, however, considered to be interplay between the destruction of elastin and collagen in the media and adventitia and apoptotic cell death of medial SMCs along with thinning of the vessel wall[48]. Some studies have reported that anti-inflammatory interventions in AAA may even accelerate disease progression[55]. However, the specific role of various cell populations of vascular origin in aneurysmal disease remains uncertain. We hypothesize that along with transmural infiltration of inflammatory cells at the early stages of the disease, SMCs contribute to the expansion of aneurysmal growth by participating in the degradation of ECM and increasing the aortic stiffness. Notch1 deficiency in SMCs may interfere with the aortic expansion initiated by the influx of inflammatory macrophages. It is not clear however, if these two factors act independently of each other or macrophages influence SMC by the release of chemokines and cytokines. Some studies have demonstrated that macrophage infiltration induces SMC apoptosis and increased MMP activity, thus contributing to aneurysmal expansion [56]. While our data and other studies have shown that SMCs are independent and actively participate in the process of degrading the aortic matrix [48, 49]. Interestingly, in humans, anti-inflammatory treatment has not been successful in preventing AAA progression making VSMCs an appealing target for further detailed study [57, 58]. Whilst synthetic VSMCs secrete ECM proteins including collagens and elastin, they also secrete MMPs that are involved in ECM breakdown activities [48, 59]. The migration of medial SMCs into intima and

media has been shown to play a decisive role in the progression of vascular diseases[<u>60</u>]. Obviously, because of the complexity of AAA pathophysiology and numerous pathways involved in the AAA formation and progression, mere decrease in the inflammatory macrophages may not be sufficient to repair pre-existing aortic dilation and elastin degradation.

Studies have shown that matricellular proteins such as CTGF play an important role in vascular remodeling by regulating ECM deposition, cell proliferation, and de-differentiation of contractile SMC to synthetic SMC phenotype[61]. In human ascending aortic aneurysms, increased reactive oxygen species (ROS) accumulation correlates with media layer degeneration and increased CTGF expression, which modulate the synthetic VSMC phenotype[50]. In addition, CTGF is also involved in monocyte migration and adhesion to injured vessel wall[50, 51]. Our findings are consistent with the previous studies and suggest that CTGF protein is overexpressed in the progression of AAA and correlates with synthetic SMCs[50, 62]. We postulate that Notch1 deficiency in the SMCs decreases the CTGF expression thus protecting against the expansion of aneurysm (Fig 7). Although the mechanism by which CTGF expression is induced in this mouse model is still obscure, possible upregulation of CTGF could be achieved by AngII or by oxidative stress in the medial layers of AAA[50]. Various fibrogenic stimuli including TGF- β , thrombin or mechanical stretch have also been reported to induce CTGF expression. Further studies may be required to define the precise mechanism by which CTGF is activated in the setting of vascular diseases including AAA.

The opposite effects of SMC-specific *Notch1* haploinsufficiency on Smad2 and Smad3 indicate that the crosstalk between Notch1 and Tgf- β signaling pathway in aneurysm may have differential effects on the downstream targets. Indeed, studies have shown that loss of *Smad3* in a mouse model drastically increases wall thickening of the abdominal aorta accompanied by vessel wall remodeling with elastin fragmentation[63]. Studies by Canalis et al[53] have shown that Notch activation directly induces *CTGF* expression in osteoblasts, and that these two pluripotent pathways may act in concert to regulate osteogenic homeostasis. It will be interesting to see if these interactions play a causal role in vascular diseases. In summary, our findings support the hypothesis that CTGF may play a role in AAA pathogenesis by promoting SMC differentiation into proliferative and apoptotic phenotype leading to aortic expansion and thus making CTGF a potential target for future therapies targeting SMC-phenotypic modulation.

Our present work demonstrates that Notch1, not Notch3 is involved in the CTGF-mediated synthetic phenotypic modulation of SMCs. Furthermore, the differential roles of Notch1 in various vascular cells observed in our study demonstrate the complexity of Notch signaling and warrant detailed future studies into the individual contributions of Notch receptors in each vascular cell type on AAA development (Fig 7). Taken together, these data suggest that Notch1 signaling has differential effects on SMC phenotype and provide strong evidence that the phenotypic modulation of SMC-phenotype can influence AAA progression. These studies might prove useful in the future for tailored cell-specific therapy and improve therapeutic efficacy to mitigate the deleterious effects of AAA progression.

Supporting information

S1 Fig

Endothelial cell (EC)-specific *Notch1* haploinsufficiency does not affect AAA development in AngII-induced mouse model.

(A) Representative images of the cross-section of the abdominal aorta of the EC-specific *Notch1* haploinsufficient *Apoe^{-/-}* mice 28 days after AngII treatment. H&E staining showing transmural

inflammatory cell infiltration; elastin staining showing fragmentation of the elastin fibers; trichrome staining showing presence of collagen fibers (blue) interspersed within the medial and adventitial layer. Survival graphs represent percentage survival in cell-specific *Notch1^{+/-};Apoe^{-/-}* as compared to *Apoe^{-/-}* mice during 28 days AngII infusion. Scale bar, 1 mm in (A), 50 µm in **B-E.**

(TIF)

Click here for additional data file.^(2.2M, TIF)

S2 Fig

SMC-specific *Notch1* haploinsufficiency does not affect dilatation of proximal aorta in AngII-induced mouse model.

Representative aortae isolated from $Apoe^{-/-}$ mice (A); $M\varphi$ -Notch1^{+/-}; $Apoe^{-/-}$ mice (B); SMC-Notch1^{+/-}; $Apoe^{-/-}$ mice (C); or EC-Notch1^{+/-}; $Apoe^{-/-}$ mice (D) treated with AngII (28 d treatment). Images were taken using Zeiss Stemi 2000-C microscope. Scale bar, 1 mm. Quantitative measurement of maximal aortic width (mm) of different groups at the ascending aorta (E), aortic arch (F), and descending aorta (G). Each symbol represents an individual animal. Mean and SEM are shown. Representative images of the cross-section of the abdominal aorta of different groups 28 d after AngII treatment with H&E staining (H-S). Scale bar, 1 mm (A-D), 50 µm in H-S.

(TIF)

Click here for additional data file.^(2.7M, TIF)

S3 Fig

Cell-specific Notch1 haploinsufficiency reduces matrix metalloproteinase production and prevents apoptotic cell death.

Immunohistochemical staining for Mmp2 (A-D), Mmp9 (E-H) and active caspase-3 (I-L) performed on cross section of abdominal aortae isolated from $Apoe^{-/-}$ and cell-specific *Notch1* haploinsufficient $Apoe^{-/-}$ mice after 28 d AngII treatment. Scale bars = 50 µm.

(TIF)

Click here for additional data file.^(4.0M, TIF)

S4 Fig

Notch1 haploinsufficiency prevents LPS-induced apoptotic cell death of SMCs.

Cells were grown at ~70% confluence, and treated with LPS (100 ng/ml for 24 h). The cells were collected, stained with annexin V-FITC (5 μ l/1x10⁵ cells suspended in 100 μ l) in binding buffer for 20 min in the dark, washed 3 times with PBS, and incubated in PI solution (5 μ l/100 μ l) immediately before 10,000 events were acquired in a Becton Dickinson LSRII flow cytometer (**A**, **B**). The percentage of annexin V-positive/PI-negative were quantified with Flow

Jo software. Bar graphs show data (mean \pm SEM) from three independent experiments (C).

(TIF)

Click here for additional data file.^(1.2M, TIF)

S5 Fig

Myeloid-specific Notch1 haploinsufficiency promotes M2-polarozation of macrophages.

Immunohistochemical staining for NICD (A-D), M1 macrophage marker, Irf8 (E-H) and Mcp1 (I-L) and M2 macrophage marker, Cd206 (M-P). Bar graphs represent fold change expression of *iNos* (Q) and *Arg1* (R) in the aorta of AngII treated *Apoe^{-/-}* and cell-specific *Notch1* haploinsufficient *Apoe^{-/-}* mice. The qPCR data were standardized to *Rpl13* and reported as ratio (mean \pm SEM, n = 3 for each group) to empty plasmids for NICD/dnMAML or non-specific siRNA. ***P<0.001, **P<0.05. Scale bars = 50 µm.

(TIF)

Click here for additional data file.^(6.5M, TIF)

S6 Fig

SMC-specific Notch1 haploinsufficiency preserves contractile-phenotype of SMCs.

Immunohistochemical staining for contractile smooth muscle cell (SMC) phenotype marker, smooth muscle-alpha-actin (SMA- α ; **A-D**) and synthetic SMC marker, vimentin (**E-H**) performed on cross section of abdominal aortae isolated from cell-specific *Notch1* haploinsufficient *Apoe^{-/-}* mice after 28 d AngII treatment. Scale bars = 50 µm.

(TIF)

Click here for additional data file. (3.0M, TIF)

S7 Fig

SMC-specific *Notch1* haploinsufficiency decreases the expression of CTGF in TGFβ-dependent manner.

(A) Western blot for Ctgf in WT and *Notch1* haploinsufficient SMCs, (B) CTGF protein expression in human aortic smooth muscle cells treated with DAPT for 24 hours followed by TGF- β agonist or inhibitor SB431542 [activin receptor-like kinase (ALK5; the TGF- β type I receptor inhibitor] treatment for 24 hours. (C) CTGF protein expression in human aortic smooth muscle cells 48 hours post transfected with Notch1 intracellular domain (NICD) overexpressing plasmid. GAPDH was used as a loading control for protein normalization in Western blotting in A-C.

(TIF)

Click here for additional data file. (102K, TIF)

S8 Fig

Macrophages from either *WT* or *Notch1*^{+/-} mice did not influence the expression of synthetic-phenotype of SMCs.

Bar graphs represent fold change in gene expression of Ctgf (A), smMHC (B), Mmp12 (C), *vimentin* (D) using mRNA obtained from *WT* SMCs co-cultured with *WT* or *Notch1*^{+/-} bone marrow derived macrophages (BMDMs) in the absence or presence of LPS/IFN- γ .

(TIF)

Click here for additional data file.(1.1M, TIF)

S1 Table

Schematic presentation to generate cell-specific Notch1 haploinsufficient mice on *Apoe^{-/-}* background.

(TIF)

Click here for additional data file.^(1.2M, TIF)

S2 Table

List of primer sequences used in genotyping (as suggested by Jackson lab).

(TIF)

Click here for additional data file. (1.5M, TIF)

S3 Table

Genotyping for cell-specific Notch1 haploinsufficient mice.

(TIF)

Click here for additional data file.^(675K, TIF)

S4 Table

List of antibodies used for immunohistochemistry and immunofluorescence.

(TIF)

Click here for additional data file. (1.0M, TIF)

S5 Table

List of primer sequences used in quantitative real-time reverse transcriptase PCR studies.

(TIF)

Click here for additional data file.^(571K, TIF)

Acknowledgments

The authors thank Dave Dunaway in the Flow Cytometry Core, members of the Morphology Core at the Research Institute at Nationwide Children's Hospital for technical support, and Yongjie Miao for the statistical analysis of the data. Most of the work related to this manuscript was performed at The Research Institute of Nationwide Children's Hospital, Columbus OH.

Funding Statement

This work was supported by R01 (HL124155) and an AHA-National Center SDG and funding from the Research Institute at Nationwide Children's Hospital to Dr. Hans.

Data Availability

All relevant data are within the paper and its Supporting Information files.

References

1. Wanhainen A. How to define an abdominal aortic aneurysm—influence on epidemiology and clinical practice. Scand J Surg. 2008;97(2):105–9. Epub 2008/06/26. doi: <u>10.1177/145749690809700204</u> [PubMed] [Google Scholar]

2. Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Blaha MJ, et al. Heart disease and stroke statistics—2014 update: a report from the American Heart Association. Circulation. 2014;129(3):e28–e292. Epub 2013/12/20. doi: <u>10.1161/01.cir.0000441139.02102.80</u> [PMC free article] [PubMed] [Google Scholar]

 Moxon JV, Parr A, Emeto TI, Walker P, Norman PE, Golledge J. Diagnosis and Monitoring of Abdominal Aortic Aneurysm: Current Status and Future Prospects. Current Problems in Cardiology. 2010;35(10):512–48. doi: <u>10.1016/j.cpcardiol.2010.08.004</u> [PMC free article] [PubMed] [Google Scholar]

4. Cooper DG, King JA, Earnshaw JJ. Role of medical intervention in slowing the growth of small abdominal aortic aneurysms. Postgraduate Medical Journal. 2009;85(1010):688–92. doi: 10.1136/pgmj.2009.085498 [PubMed] [Google Scholar]

5. Hurks R, Hoefer IE, Vink A, Pasterkamp G, Schoneveld A, Kerver M, et al. Different Effects of Commonly Prescribed Statins on Abdominal Aortic Aneurysm Wall Biology. European Journal of Vascular and Endovascular Surgery. 2010;39(5):569–76. doi: <u>10.1016/j.ejvs.2010.01.030</u> [PubMed] [Google Scholar]

6. Baxter BT, Terrin MC, Dalman RL. Medical Management of Small Abdominal Aortic Aneurysms. Circulation. 2008;117(14):1883–9. doi: <u>10.1161/CIRCULATIONAHA.107.735274</u> [PMC free article] [PubMed] [Google Scholar]

7. Curci JA. Digging in the "Soil" of the Aorta to Understand the Growth of Abdominal Aortic Aneurysms. Vascular. 2009;17(suppl 1):S21–S9. [PMC free article] [PubMed] [Google Scholar]

8. Lu H, Rateri D, Bruemmer D, Cassis L, Daugherty A. Novel Mechanisms of Abdominal Aortic Aneurysms. Curr Atheroscler Rep. 2012;14(5):402–12. doi: <u>10.1007/s11883-012-0271-y</u>

[PMC free article] [PubMed] [Google Scholar]

9. Uchida HA, Kristo F, Rateri DL, Lu H, Charnigo R, Cassis LA, et al. Total lymphocyte deficiency attenuates AngII-induced atherosclerosis in males but not abdominal aortic aneurysms in apoE deficient mice. Atherosclerosis. 2010;211(2):399–403. doi: <u>10.1016/j.atherosclerosis.2010.02.034</u> [PMC free article] [PubMed] [Google Scholar]

10. Saraff K, Babamusta F, Cassis LA, Daugherty A. Aortic Dissection Precedes Formation of Aneurysms and Atherosclerosis in Angiotensin II-Infused, Apolipoprotein E-Deficient Mice. Arteriosclerosis, Thrombosis, and Vascular Biology. 2003;23(9):1621–6. doi: 10.1161/01.ATV.0000085631.76095.64 [PubMed] [Google Scholar]

11. Reeps C, Pelisek J, Seidl S, Schuster T, Zimmermann A, Kuehnl A, et al. Inflammatory infiltrates and neovessels are relevant sources of MMPs in abdominal aortic aneurysm wall. Pathobiology: journal of immunopathology, molecular and cellular biology. 2009;76(5):243–52. Epub 2009/10/10. [PubMed] [Google Scholar]

12. Raines EW, Ross R. Smooth muscle cells and the pathogenesis of the lesions of atherosclerosis. British Heart Journal. 1993;69(1 Suppl):S30–S7. [PMC free article] [PubMed] [Google Scholar]

13. Wamhoff BR, Hoofnagle MH, Burns A, Sinha S, McDonald OG, Owens GK. A G/C element mediates repression of the SM22alpha promoter within phenotypically modulated smooth muscle cells in experimental atherosclerosis. Circulation research. 2004;95(10):981–8. Epub 2004/10/16. doi: 10.1161/01.RES.0000147961.09840.fb [PubMed] [Google Scholar]

14. Yoshida T, Kawai-Kowase K, Owens GK. Forced expression of myocardin is not sufficient for induction of smooth muscle differentiation in multipotential embryonic cells. Arterioscler Thromb Vasc Biol. 2004;24(9):1596–601. Epub 2004/07/03. doi: <u>10.1161/01.ATV.0000137190.63214.c5</u> [PubMed] [Google Scholar]

15. Cheng P, Gabrilovich D. Notch signaling in differentiation and function of dendritic cells. Immunol Res. 2008;41(1):1–14. Epub 2007/11/17. PubMed Central PMCID: PMC2709100. doi: 10.1007/s12026-007-8011-z [PMC free article] [PubMed] [Google Scholar]

16. Monsalve E, Perez MA, Rubio A, Ruiz-Hidalgo MJ, Baladron V, Garcia-Ramirez JJ, et al. Notch-1 up-regulation and signaling following macrophage activation modulates gene expression patterns known to affect antigen-presenting capacity and cytotoxic activity. J Immunol. 2006;176(9):5362–73. Epub 2006/04/20. [PubMed] [Google Scholar]

17. Ohishi K, Varnum-Finney B, Serda RE, Anasetti C, Bernstein ID. The Notch ligand, Delta-1, inhibits the differentiation of monocytes into macrophages but permits their differentiation into dendritic cells. Blood. 2001;98(5):1402–7. Epub 2001/08/25. [PubMed] [Google Scholar]

Gridley T. Notch signaling in vascular development and physiology. Development.
 2007;134(15):2709–18. Epub 2007/07/06. doi: <u>10.1242/dev.004184</u> [PubMed] [Google Scholar]

19. Iso T, Hamamori Y, Kedes L. Notch signaling in vascular development. Arterioscler Thromb Vasc Biol. 2003;23(4):543–53. Epub 2003/03/05. doi: <u>10.1161/01.ATV.0000060892.81529.8F</u> [PubMed] [Google Scholar]

20. Lawson ND, Scheer N, Pham VN, Kim CH, Chitnis AB, Campos-Ortega JA, et al. Notch signaling is required for arterial-venous differentiation during embryonic vascular development. Development. 2001;128(19):3675–83. Epub 2001/10/05. [PubMed] [Google Scholar]

21. Gridley T. Notch signaling during vascular development. Proc Natl Acad Sci U S A.

2001;98(10):5377–8. Epub 2001/05/10. PubMed Central PMCID: PMC33218. doi: 10.1073/pnas.101138098 [PMC free article] [PubMed] [Google Scholar]

22. Gasperowicz M, Otto F. The Notch Signalling Pathway in the Development of the Mouse Placenta. Placenta. 2008;29(8):651–9. doi: <u>10.1016/j.placenta.2008.06.004</u> [PubMed] [Google Scholar]

23. Talora C, Campese AF, Bellavia D, Felli MP, Vacca A, Gulino A, et al. Notch signaling and diseases: An evolutionary journey from a simple beginning to complex outcomes. Biochimica et Biophysica Acta (BBA)—Molecular Basis of Disease. 2008;1782(9):489–97. [PubMed] [Google Scholar]

24. Quillard T, Charreau B. Impact of Notch Signaling on Inflammatory Responses in Cardiovascular Disorders. International Journal of Molecular Sciences. 2013;14(4):6863–88. doi: 10.3390/ijms14046863 [PMC free article] [PubMed] [Google Scholar]

25. Sweeney C, Morrow D, Birney YA, Coyle S, Hennessy C, Scheller A, et al. Notch 1 and 3 receptors modulate vascular smooth muscle cell growth, apoptosis and migration via a CBF-1/RBP-Jk dependent pathway. The FASEB Journal. 2004. [PubMed] [Google Scholar]

26. Rostama B, Peterson SM, Vary CPH, Liaw L. Notch signal integration in the vasculature during remodeling. Vascular Pharmacology. 2014;63(2):97–104. doi: <u>10.1016/j.vph.2014.10.003</u> [PMC free article] [PubMed] [Google Scholar]

27. Zheng CG, Chen R, Xie JB, Liu CB, Jin Z, Jin C. Immunohistochemical expression of Notch1, Jagged1, NF-kappaB and MMP-9 in colorectal cancer patients and the relationship to clinicopathological parameters. Cancer biomarkers: section A of Disease markers. 2015;15(6):889–97. Epub 2015/09/26. doi: 10.3233/CBM-150533 [PubMed] [Google Scholar]

28. Wang Y-W, Ren H-L, Wang H-F, Li F-D, Li H-H, Zheng Y-H. Combining detection of Notch1 and tumor necrosis factor-α converting enzyme is a reliable biomarker for the diagnosis of abdominal aortic aneurysms. Life Sciences. 2015;127:39–45. doi: 10.1016/j.lfs.2015.02.009 [PubMed] [Google Scholar]

29. Kuivaniemi H, Ryer EJ, Elmore JR, Tromp G. Understanding the pathogenesis of abdominal aortic aneurysms. Expert Rev Cardiovasc Ther. 2015;13(9):975–87. PubMed Central PMCID: PMCPMC4829576. doi: <u>10.1586/14779072.2015.1074861 [PMC free article]</u> [PubMed] [Google Scholar]

30. Gridley T. Notch signaling in the vasculature. Current topics in developmental biology. 2010;92:277–309. doi: <u>10.1016/S0070-2153(10)92009-7</u> [PMC free article] [PubMed] [Google Scholar]

31. Tanigaki K, Honjo T. Chapter Seven—Two Opposing Roles of RBP-J in Notch Signaling In: Raphael K, editor. Current topics in developmental biology. Volume 92: Academic Press; 2010. p. 231–52. [PubMed] [Google Scholar]

32. Cheng J, Koenig SN, Kuivaniemi HS, Garg V, Hans CP. Pharmacological inhibitor of notch signaling stabilizes the progression of small abdominal aortic aneurysm in a mouse model. J Am Heart Assoc. 2014;3(6):e001064 PubMed Central PMCID: PMCPMC4338693. doi: 10.1161/JAHA.114.001064 [PMC free article] [PubMed] [Google Scholar]

33. Hans CP, Koenig SN, Huang N, Cheng J, Beceiro S, Guggilam A, et al. Inhibition of Notch1 signaling reduces abdominal aortic aneurysm in mice by attenuating macrophage-mediated inflammation. Arterioscler Thromb Vasc Biol. 2012;32(12):3012–23. Epub 2012/10/23. doi: 10.1161/ATVBAHA.112.254219 [PMC free article] [PubMed] [Google Scholar]

34. Hans CP, Zerfaoui M, Naura AS, Catling A, Boulares AH. Differential effects of PARP inhibition on vascular cell survival and ACAT-1 expression favouring atherosclerotic plaque stability. Cardiovasc Res. 2008;78(3):429–39. doi: 10.1093/cvr/cvn018 [PubMed] [Google Scholar]

35. Oumouna-Benachour K, Hans CP, Suzuki Y, Naura A, Datta R, Belmadani S, et al. Poly(ADPribose) polymerase inhibition reduces atherosclerotic plaque size and promotes factors of plaque stability in apolipoprotein E-deficient mice: effects on macrophage recruitment, nuclear factor-kappaB nuclear translocation, and foam cell death. Circulation. 2007;115(18):2442–50. doi: 10.1161/CIRCULATIONAHA.106.668756 [PubMed] [Google Scholar]

36. Zerfaoui M, Suzuki Y, Naura AS, Hans CP, Nichols C, Boulares AH. Nuclear translocation of p65 NF-kappaB is sufficient for VCAM-1, but not ICAM-1, expression in TNF-stimulated smooth muscle cells: Differential requirement for PARP-1 expression and interaction. Cell Signal. 2008;20(1):186–94. doi: <u>10.1016/j.cellsig.2007.10.007</u> [PMC free article] [PubMed] [Google Scholar]

37. Acharya A, Hans CP, Koenig SN, Nichols HA, Galindo CL, Garner HR, et al. Inhibitory role of Notch1 in calcific aortic valve disease. PloS one. 2011;6(11):e27743 Epub 2011/11/24. PubMed Central PMCID: PMC3218038. doi: <u>10.1371/journal.pone.0027743</u> [PMC free article] [PubMed] [Google Scholar]

38. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an opensource platform for biological-image analysis. Nat Meth. 2012;9(7):676–82. [PMC free article] [PubMed] [Google Scholar]

39. Bosse K, Hans CP, Zhao N, Koenig SN, Huang N, Guggilam A, et al. Endothelial nitric oxide signaling regulates Notch1 in aortic valve disease. Journal of Molecular and Cellular Cardiology. 2013;60(0):27–35. [PMC free article] [PubMed] [Google Scholar]

40. Lin C-H, Lilly B. Endothelial Cells Direct Mesenchymal Stem Cells Toward a Smooth Muscle Cell Fate. Stem cells and development. 2014;23(21):2581–90. doi: <u>10.1089/scd.2014.0163</u> [<u>PMC free article</u>] [<u>PubMed</u>] [<u>Google Scholar</u>]

41. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. Nat Meth. 2012;9(7):671–5. [PMC free article] [PubMed] [Google Scholar]

42. Liu H, Zhang W, Kennard S, Caldwell RB, Lilly B. Notch3 is critical for proper angiogenesis and mural cell investment. Circulation research. 2010;107(7):860–70. doi: 10.1161/CIRCRESAHA.110.218271 [PMC free article] [PubMed] [Google Scholar]

43. Baker M, Robinson SD, Lechertier T, Barber PR, Tavora B, D'Amico G, et al. Use of the mouse aortic ring assay to study angiogenesis. Nat Protoc. 2012;7(1):89–104. [PubMed] [Google Scholar]

44. Koenig SN, Bosse KM, Nadorlik HA, Lilly B, Garg V. Evidence of Aortopathy in Mice with Haploinsufficiency of Notch1 in Nos3-Null Background. Journal of cardiovascular development and disease. 2015;2(1):17–30. doi: <u>10.3390/jcdd2010017</u> [PMC free article] [PubMed] [Google Scholar]

45. Guo D-C, Papke CL, He R, Milewicz DM. Pathogenesis of Thoracic and Abdominal Aortic Aneurysms. Annals of the New York Academy of Sciences. 2006;1085(1):339–52. [PubMed] [Google Scholar]

46. McKellar SH, Tester DJ, Yagubyan M, Majumdar R, Ackerman MJ, Sundt TM 3rd. Novel NOTCH1 mutations in patients with bicuspid aortic valve disease and thoracic aortic aneurysms. J Thorac Cardiovasc Surg. 2007;134(2):290–6. Epub 2007/07/31. doi: <u>10.1016/j.jtcvs.2007.02.041</u> [PubMed] [Google Scholar] 47. Zou S, Ren P, Nguyen M, Coselli JS, Shen YH, LeMaire SA. Notch Signaling in Descending Thoracic Aortic Aneurysm and Dissection. PloS one. 2012;7(12):e52833 doi: <u>10.1371/journal.pone.0052833</u> [PMC free article] [PubMed] [Google Scholar]

48. Airhart N, Brownstein BH, Cobb JP, Schierding W, Arif B, Ennis TL, et al. Smooth muscle cells from abdominal aortic aneurysms are unique and can independently and synergistically degrade insoluble elastin. Journal of Vascular Surgery. 2014;60(4):1033–42.e5. doi: <u>10.1016/j.jvs.2013.07.097</u> [PMC free article] [PubMed] [Google Scholar]

49. Ailawadi G, Moehle CW, Pei H, Walton SP, Yang Z, Kron IL, et al. Smooth muscle phenotypic modulation is an early event in aortic aneurysms. The Journal of Thoracic and Cardiovascular Surgery. 2009;138(6):1392–9. doi: 10.1016/j.jtcvs.2009.07.075 [PMC free article] [PubMed] [Google Scholar]

50. Branchetti E, Poggio P, Sainger R, Shang E, Grau JB, Jackson BM, et al. Oxidative stress modulates vascular smooth muscle cell phenotype via CTGF in thoracic aortic aneurysm. Cardiovasc Res. 2013;100(2):316–24. Epub 2013/08/30. PubMed Central PMCID: PMC4192047. doi: <u>10.1093/cvr</u>/<u>/cvt205 [PMC free article] [PubMed] [Google Scholar]</u>

51. Cicha I, Zitzmann R, Goppelt-Struebe M. Dual inhibition of Src family kinases and Aurora kinases by SU6656 modulates CTGF (connective tissue growth factor) expression in an ERK-dependent manner. Int J Biochem Cell Biol. 2014;46:39–48. Epub 2013/11/28. doi: <u>10.1016/j.biocel.2013.11.014</u> [PubMed] [Google Scholar]

52. Ha DM, Carpenter LC, Koutakis P, Swanson SA, Zhu Z, Hanna M, et al. Transforming growth factor-beta 1 produced by vascular smooth muscle cells predicts fibrosis in the gastrocnemius of patients with peripheral artery disease. Journal of Translational Medicine. 2016;14:39 doi: 10.1186/s12967-016-0790-3 [PMC free article] [PubMed] [Google Scholar]

53. Canalis E, Zanotti S, Smerdel-Ramoya A. Connective Tissue Growth Factor is a Target of Notch Signaling in Cells of the Osteoblastic Lineage. Bone. 2014;64:273–80. doi: <u>10.1016/j.bone.2014.04.028</u> [<u>PMC free article</u>] [<u>PubMed</u>] [<u>Google Scholar</u>]

54. Daugherty A, Manning M, Cassis L. Angiotensin II promotes atherosclerotic lesions and aneurysms in apolipoprotein E-deficient mice. J Clin Invest. 2000;105:1605–12. doi: <u>10.1172/JCI7818</u> [PMC free article] [PubMed] [Google Scholar]

55. Lindeman JHN, Rabelink TJ, van Bockel JH. Immunosuppression and the Abdominal Aortic Aneurysm: Doctor Jekyll or Mister Hyde? Circulation. 2011;124(18):e463–e5. doi: 10.1161/CIRCULATIONAHA.110.008573 [PubMed] [Google Scholar]

56. Wang Q, Ren J, Morgan S, Liu Z, Dou C, Liu B. Monocyte Chemoattractant Protein-1 (MCP-1) Regulates Macrophage Cytotoxicity in Abdominal Aortic Aneurysm. PloS one. 2014;9(3):e92053 doi: <u>10.1371/journal.pone.0092053 [PMC free article] [PubMed] [Google Scholar]</u>

57. Englesbe MJ, Wu AH, Clowes AW, Zierler RE. The prevalence and natural history of aortic aneurysms in heart and abdominal organ transplant patients. J Vasc Surg. 2003;37(1):27–31. Epub 2003/01/07. doi: <u>10.1067/mva.2003.57</u> [PubMed] [Google Scholar]

58. Wilson WRW, Wills J, Furness PN, Loftus IM, Thompson MM. Abdominal Aortic Aneurysm Rupture is not Associated with an Up-regulation of Inflammation within the Aneurysm Wall. European Journal of Vascular and Endovascular Surgery. 2010;40(2):191–5. doi: <u>10.1016/j.ejvs.2010.04.014</u> [PubMed] [Google Scholar]

59. Liu S-L, Bae YH, Yu C, Monslow J, Hawthorne EA, Castagnino P, et al. Matrix metalloproteinase-12 is an essential mediator of acute and chronic arterial stiffening. Scientific Reports.

2015;5:17189 doi: 10.1038/srep17189 [PMC free article] [PubMed] [Google Scholar]

60. Akyurek LM, Paul LC, Funa K, Larsson E, Fellstrom BC. Smooth muscle cell migration into intima and adventitia during development of transplant vasculopathy. Transplantation. 1996;62(10):1526–9. [PubMed] [Google Scholar]

61. Beamish JA, He P, Kottke-Marchant K, Marchant RE. Molecular Regulation of Contractile Smooth Muscle Cell Phenotype: Implications for Vascular Tissue Engineering. Tissue Engineering Part B, Reviews. 2010;16(5):467–91. doi: <u>10.1089/ten.TEB.2009.0630</u> [PMC free article] [PubMed] [Google Scholar]

62. Gu J, Liu X, Wang Q-x, Tan H-w, Guo M, Jiang W-f, et al. Angiotensin II increases CTGF expression via MAPKs/TGF-β1/TRAF6 pathway in atrial fibroblasts. Experimental cell research. 2012;318(16):2105–15. doi: 10.1016/j.yexcr.2012.06.015 [PubMed] [Google Scholar]

63. Dai X, Shen J, Priyanka Annam N, Jiang H, Levi E, Schworer CM, et al. SMAD3 deficiency promotes vessel wall remodeling, collagen fiber reorganization and leukocyte infiltration in an inflammatory abdominal aortic aneurysm mouse model. Scientific Reports. 2015;5:10180 doi: 10.1038/srep10180 [PMC free article] [PubMed] [Google Scholar]

Articles from PLoS ONE are provided here courtesy of Public Library of Science